

X-RAY, OPTICAL, AND INFRARED IMAGING AND SPECTRAL PROPERTIES OF THE 1 MS *CHANDRA* DEEP FIELD NORTH SOURCES

A. J. BARGER,^{1,2,3} L. L. COWIE,³ W. N. BRANDT,⁴ P. CAPAK,³ G. P. GARMIRE,⁴
 A. E. HORNSCHMEIER,⁴ A. T. STEFFEN,¹ E. H. WEHNER¹

Submitted to the Astronomical Journal

ABSTRACT

We present the optical, near-infrared, submillimeter, and radio follow-up catalog of the X-ray selected sources from an ≈ 1 Ms *Chandra* observation of the Hubble Deep Field North region. We have B , V , R , I , and z' magnitudes for the 370 X-ray point sources, HK' magnitudes for 276, and spectroscopic redshifts for 182. We present high-quality spectra for 175 of these. The redshift distribution shows indications of structures at $z = 0.843$ and $z = 1.0175$ (also detected in optical surveys) which could account for a part of the field-to-field variation seen in the X-ray number counts; however, these structures do not dominate the number of X-ray sources in the sample and hence should not strongly affect the redshift distribution. All of the X-ray sources with $z > 1.6$ are either broad-line AGN or have narrow Ly α and/or CIII] 1909 Å emission; none of the known $z > 1.6$ absorption-line galaxies in the field are detected individually in X-rays. We estimate photometric redshifts for the sources with $B - I > 1.5$ (bluer than this it is hard to distinguish between low redshift irregulars and luminous high-redshift AGN) and find agreement (most are within 25%) with the available spectroscopic redshifts. The majority of the galaxies in both the 2 – 8 keV (hard) and 0.5 – 2 keV (soft) samples have absolute magnitudes comparable to or more luminous than $M_I^* = -22$. The flux contributions separated into unit bins of redshift show that the $z < 1$ spectroscopically identified sources already contribute about one-third of the total flux in both the hard and soft bands. Thus, major accretion onto supermassive black holes has occurred since the Universe was half its present age. We find from ratios of the X-ray counts that the X-ray spectra are well-described by absorption of an intrinsic $\Gamma = 1.8$ power-law, with N_H values ranging from about 10^{21} cm⁻² to 5×10^{23} cm⁻². We find very little evolution in the maximum rest-frame opacity-corrected and K -corrected 2 – 8 keV X-ray luminosities with decreasing redshift until $z \lesssim 0.5$, where the volume becomes too small to probe effectively very high luminosity sources. We estimate that the *Chandra* sources that produce 87% of the *HEAO-A* X-ray background (XRB) at 3 keV produce 57% at 20 keV, provided that at high energies the spectral shape of the sources continues to be well-described by a $\Gamma = 1.8$ power-law. However, when the *Chandra* contributions are renormalized to the *BeppoSAX* XRB at 3 keV, the shape matches fairly well the observed XRB at both energies. Thus, whether a substantial population of as-yet undetected Compton-thick sources, or some change in the spectral shape of the current sources from the simple power-law dependence, is required to completely resolve the XRB above 10 keV depends critically on how the currently discrepant XRB measurements in the 1 – 10 keV energy range tie together with the higher energy XRB.

Subject headings: cosmology: observations — galaxies: evolution — galaxies: formation

1. INTRODUCTION

X-ray surveys most directly trace accretion onto supermassive black holes and hence provide our best window on black hole evolution. Some primary observational goals of X-ray studies include the measurement of supermassive black hole properties (such as number density and accretion rates) versus redshift and the determination of the nature of the host galaxies. Since much of the accretion power in the Universe may be absorbed by substantial neutral hydrogen column densities (e.g., Fabian & Iwasawa 1999), the challenge is to construct a *complete census* of supermassive black holes to the earliest epoch, including sources which are heavily obscured from soft X-ray energies to the near-infrared (NIR). The census is well underway for unobscured sources (i.e., quasars with broad

emission lines and big blue bumps), but the obscured population is not well-characterized, except for the relatively rare radio galaxies.

A fundamental goal of the *Chandra X-ray Observatory* was to resolve the hard (2 – 8 keV) X-ray background (XRB) into discrete sources. At these energies the photons can penetrate all but the highest column densities ($> 10^{24}$ cm⁻²) of gas and dust, so many obscured active galactic nuclei (AGN) can now be detected. The two recent ≈ 1 Ms exposures of the *Chandra* Deep Field North (CDF-N; Brandt et al. 2001c, hereafter B01) and the *Chandra* Deep Field South (CDF-S; Giacconi et al. 2002) have resolved $> 80 - 90\%$ of the 2 – 8 keV XRB into discrete sources (Campana et al. 2001; Cowie et al. 2002; Rosati et al. 2002) with the main uncertainty being the normalization of the XRB itself. The CDF-N exposure

¹Department of Astronomy, University of Wisconsin-Madison, 475 North Charter Street, Madison, WI 53706

²Department of Physics and Astronomy, University of Hawaii, 2505 Correa Road, Honolulu, HI 96822

³Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

⁴Department of Astronomy & Astrophysics, 525 Davey Laboratory, The Pennsylvania State University, University Park, PA 16802

has recently been extended to a second megasecond.

Although with the *Chandra* and *XMM-Newton Observatories* there has been rapid improvement in our ability to detect hard X-ray sources at the faintest fluxes, our efforts to understand in detail the nature and evolution of the sources creating the hard XRB are still in the early stages. Optical and NIR follow-up studies of 100–300 ks *Chandra* (Mushotzky et al. 2000; Hornschemeier et al. 2001, hereafter H01; Barger et al. 2001a, c; Tozzi et al. 2002; Stern et al. 2002b) and *XMM-Newton* (Hasinger et al. 2001; Baldi et al. 2002; Fiore et al. 2001) imaging surveys have found that almost half of the hard X-ray light arises in optically bright ($I < 23.5$) galaxies in the $z < 1.5$ redshift range. Almost all of these sources can be spectroscopically identified, and most are bulge-dominated galaxies with near- L^* luminosities. The other half of the hard X-ray light arises in a mixture of $z > 1.5$ AGN and optically faint galaxies ($I > 23.5$) that cannot be spectroscopically identified.

Contrary to the situation for the faint *ROSAT* soft X-ray sources (Hasinger et al. 1998; Schmidt et al. 1998), the vast majority ($> 80\%$) of the spectroscopically identified hard X-ray sources do not have broad optical or ultraviolet lines, and almost half show no obvious high-ionization signatures of AGN activity in their spectra. The hard to soft X-ray flux ratios of the latter sources suggest that the sources are highly absorbed systems whose high column densities could effectively extinguish the optical, ultraviolet, and NIR continua from the AGN and render traditional identification techniques ineffective.

Based on the colors, many of the spectroscopically unidentified $I > 23.5$ galaxies may lie at redshifts in the range $z = 1.5$ to 3 (Crawford et al. 2001; Cowie et al. 2001; Barger et al. 2001a, b; Alexander et al. 2001). However, a very intriguing possibility suggested by Haiman & Loeb (1999) is that some of the optically faint sources may instead be low luminosity quasars at very high redshifts ($z > 5$), since cold dark matter models can allow a large number of such high-redshift AGN.

The ≈ 1 Ms *Chandra* exposures have also resolved $> 90\%$ of the soft (0.5–2) XRB into discrete sources (B01; Rosati et al. 2002). At the faintest X-ray fluxes probed by these exposures, galaxies whose X-ray light is dominated by processes related to star formation rather than AGN activity begin to be significant contributors to the number counts, even though they contribute only 2–3% to the X-ray light (e.g., Brandt et al. 2001a; Hornschemeier et al. 2002). The large numbers of detections are understandable since we now have the sensitivity to detect the most luminous ($> 2 \times 10^{39}$ erg s $^{-1}$) X-ray binaries and supernova remnants at $z < 0.15$. AGN may cease to dominate the X-ray number counts at extremely faint fluxes (e.g., Moran, Lehnert, & Helfand 1999) in a manner analogous to the dominance of star forming galaxies rather than AGN at very faint radio fluxes (e.g., Richards 2000).

In this paper we present follow-up optical, NIR, and submillimeter imaging and optical spectroscopy of the X-ray sources detected in the ≈ 1 Ms CDF-N exposure to characterize the properties of the faint X-ray source populations. We then use our data to describe the evolution of the X-ray sources and their host galaxies.

2. X-RAY SAMPLE

B01 presented the 975.3 ks CDF-N X-ray images, along with details of the observations, data reduction, and technical analysis. B01 merged their source lists in four standard X-ray bands, 0.5–8 keV (full band), 0.5–2.0 keV (soft band), 2–8 keV (hard band), and 4–8 keV (ultrahard band), into a catalog (Table 3 in B01) of 370 significantly detected point sources over an area of about 450 arcmin 2 . Near the aim point the data reach limiting fluxes of $\approx 3 \times 10^{-17}$ (0.5–2 keV) and $\approx 2 \times 10^{-16}$ erg cm $^{-2}$ s $^{-1}$ (2–8 keV), which are, respectively, ≈ 40 and ≈ 400 times fainter than achieved by pre-*Chandra* missions. The number counts and completeness levels are described in B01 and Cowie et al. (2002). The six extended X-ray sources detected in the CDF-N are discussed in great detail in Bauer et al. (2002) and will not be considered in the present work.

In this paper we present a follow-up data table to match the full B01 X-ray point source catalog. However, outside a 10' radius from the approximate *Chandra* image center J2000 RA 12 h 36 m 51.20 s and Dec 62 $^\circ$ 13' 43.0'' (this choice of center maximizes the average exposure time within a 6.5' radius circle), the X-ray point spread function degrades rapidly; thus, in order to have a more uniform flux sample with accurate positions for our analysis, in § 9 we introduce a restricted 10' radius sample that we will use (in addition to some even more restricted subsamples) subsequently.

Table 1 lists the full 370 B01 X-ray sources by number (column 1) and by RA and Dec coordinates in decimal degrees (columns 2 and 3). Columns 4–15, to be discussed in subsequent subsections, give submillimeter fluxes, radio fluxes, isophotal R magnitudes, corrected aperture HK' , z' , I , R , V , and B magnitudes, optical Δx and Δy offsets in arcseconds from the X-ray positions, and redshifts. The final column contains a 'B' if the spectrum has broad emission lines and a 'C' if the source is optically compact.

Figure 1 shows a schematic of the 370 X-ray point sources in the full B01 sample (solid squares denote sources with spectroscopic identifications, as discussed in § 9). Radii of 6.5' and 10' from the approximate X-ray image center are indicated with large circles and will be discussed in subsequent sections. The dashed contour shows the area covered by the HK' imaging data (see § 4) and the solid contour shows the area covered by the submillimeter data (see § 6). The optical data cover a 24' $\times$ 24' area centered on the Hubble Deep Field North (HDF-N).

3. OPTICAL OBSERVATIONS AND REDUCTION

The optical imaging data consist of Johnson B , Johnson V , Cousins R , Cousins I , and Sloan z' observations obtained with the Subaru Prime Focus Camera (Suprime-Cam; Miyazaki et al. 1998) on the Subaru⁵ 8.2 m telescope between 2001 February and April. The images and catalogs of the entire sample of galaxies and stars in the field are presented in P. Capak et al., in preparation, where a more extensive discussion may be found. Here we briefly summarize the Suprime-Cam observations and reductions.

A five step dither pattern with 1' steps was used to fill in the chip gaps and provide adequate offsets for flat-fielding each dither pattern separately. Half the dither patterns

⁵The Subaru telescope is operated by the National Astronomical Observatory of Japan.

were taken at a 90 degree rotation to provide better chip-to-chip photometric calibration and remove bleeding from saturated stars. The total exposure times were 6000 s in B and V , 19440 s in R , 4080 s in I , and 11450 s in z' . Seeing was good throughout the observations, ranging from $0.5''$ to $1.5''$. The median seeing in the final images is $0.75''$ in B , $1''$ in V , $1''$ in R , $0.8''$ in I , and $0.8''$ in z' . The inverse second moment of the light in the highest resolution B image was used to determine the compactness (Kron 1980). $B < 26.5$ sources consistent with being point-like and unsaturated are marked ‘C’ for compact in the final column of Table 1.

The data were reduced using Nick Kaiser’s Imcat software package. The data were overscan corrected and bias subtracted. Bad pixels were masked. No dark subtraction was performed since Suprime-Cam has a negligible dark current. A median sky flat was generated for each band to correct for the vignetting pattern. For the I and z' bands a surface was tessellated over a grid spaced at 128 pixel intervals. This surface was used instead of a median sky flat to remove high frequency fringing, which is additive, from the flat. To generate a fringe frame, a median sky flat was created for each night in I and z' and divided by the tessellated surface. The fringe frame was scaled to the background level in each chip and subtracted to remove the fringing. Next, a separate median flat was generated for each dither pattern to remove flattening variations caused by flexure in the instrument. A second order polynomial was then fit to the sky background and subtracted to remove the sky. After this reduction the images were flat to better than 1% r.m.s.

The astrometric solution was obtained in two stages. Stars brighter than 19th magnitude are saturated in most of the science images making comparisons to the USNOA-2 difficult. Thus, a standard star frame was fit to the USNOA-2 to find a rough initial solution for each chip in the mosaic. This initial solution was applied to the science frames, and stars appearing within $2''$ of the same position in at least four images were identified. A cubic polynomial distortion was then fit to each chip of the mosaic in each exposure to minimize the r.m.s. scatter in the position of the stars from exposure to exposure. The solution was iterated several times to eliminate mismatched stars. The absolute positioning was then chosen to match the VLA 20 cm catalog of Richards (2000; see § 7), which was also used in B01 to determine the absolute positions of the *Chandra* sources.

We show z' -band thumbnail images of the CDF-N sources in Fig. 2 ordered by B01 catalog number. We use the z' -band since many of the sources are red and hence more easily seen in this band. Each thumbnail is $18''$ on a side. The number of the source from Table 1 is printed in the upper left corner. If the source is also a submillimeter source (see § 6), the letter ‘S’ is printed in the upper right corner. If the source is also a radio source (see § 7), the letter ‘R’ is printed in the upper right corner. If the source is in the B01 2 – 8 keV sample, the letter ‘H’ is printed in the bottom right corner. If there is a spectroscopic identification for a source that lies within $2''$ of the X-ray position (see § 9), then the redshift (or the word ‘STAR’ for a stellar spectrum) is printed in the bottom left corner. If there is a spectroscopic identification for a source that lies outside of our chosen cross-identification radius of $2''$

but still within $5''$ of the X-ray position, then the redshift of that source is printed in fainter text to indicate that the redshift has been given for interest. However, apart from two special cases discussed in § 9, such sources have not been cross-identified in Table 1 with the X-ray sources, nor have the redshifts been used in any subsequent analysis.

Of the 370 sources in the CDF-N catalog, 210 have measured R magnitudes brighter than 24. Since the surface density of galaxies and stars in this region is roughly 7.3 arcmin^{-2} to this magnitude limit, the number of random overlaps should be small: 5 chance projections within a $1.5''$ radius and 22 within a $3''$ radius. For the $R < 24$ sources the median offset between the optical and X-ray positions is $-0.09''$ in RA and $-0.05''$ in Dec, illustrating that the absolute astrometric solutions are indeed consistent.

The positional uncertainties are expected primarily to reflect the centroiding accuracies and the overall distortions in the X-ray data, both of which increase as we move to larger angles away from the optical axis of the *Chandra* telescope and also depend on the flux of the source (B01). In order to determine the relative positional accuracy of the X-ray and optical data, we measured the spread in offsets between the X-ray and optical positions for the $R \leq 24$ sample. Within a $6.5'$ radius from the approximate X-ray image center the r.m.s. dispersion is $0.28''$ in RA and $0.25''$ in Dec, while between $6.5'$ and $10'$ this rises to $0.58''$ in RA and $0.7''$ in Dec. The corresponding two dimensional dispersions are $0.38''$ and $0.9''$. The circles in Fig. 2 show $2''$ (5σ) radii for sources within $6.5'$ of the center and $3.6''$ radii (4σ) for sources beyond this radius. Nearly all the true counterparts should lie within these error circles.

The Suprime-Cam data were photometrically calibrated using sources in the HDF-N proper, as given in the catalog of Fernández-Soto, Lanzetta, & Yahil (1999). To facilitate comparison with previous work, we give the B and V magnitudes in the Johnson system and the R and I magnitudes in the Kron-Cousins system. The offset between AB magnitude and Vega-based magnitude is 0.41, 0.20, 0.0, and -0.11 for I , R , V , and B , respectively, where the AB magnitude is defined by $-48.60 - 2.5 \log f_\nu$. Here f_ν is the flux of the source in units of $\text{erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1}$. The z' magnitudes are given in the AB system.

We measured the magnitude of each source in a $3''$ diameter aperture. We then corrected to an approximate total magnitude using an offset calculated as the median difference between $3''$ and $6''$ aperture magnitudes measured for a complete $I < 25$ optical sample in the field. Higher resolution images were smoothed to match the seeing in the poorer resolution images in order to give the most precise color measurements. For sources with a counterpart brighter than $z' = 24$ within $2''$ of the X-ray position, we centered on the brightest z' -band peak within the $2''$ error circle. The optical Δx and Δy offsets in arcseconds from the X-ray position (positive values are to the East and North) are given in columns 13 and 14 of Table 1. For optically fainter sources we centered on the X-ray position.

For sources brighter than $R = 23$ we also measured the isophotal magnitude in the R -band to an isophotal level which is 1% of the peak surface brightness. At magnitudes fainter than $R = 20$ the isophotal and corrected aperture magnitudes generally agree well, except for the

small number of X-ray sources which lie on the edges of bright galaxies (sources 30, 62, 125, 192, 218, 225, 271, and 290). Here the isophotal magnitude measures the magnitude of the bright galaxy. At $R < 20$ the sizes of many of the galaxies become larger than the $3''$ diameter aperture, and the isophotal magnitudes become brighter than the corrected aperture magnitudes. For these objects the isophotal magnitude may be preferred.

The isophotal R magnitudes are given in column 6 of Table 1 and the corrected aperture magnitudes for z' , I , R , V , and B are given in columns 8–12. Sources which are saturated in a given band are listed at a nominal magnitude of 18 (given in brackets), which is about a magnitude brighter than where saturation occurs in the images, while sources which are contaminated by bright neighbors or by bleeding columns from bright stars are listed at a nominal magnitude of 30. Sources with a negative flux in the aperture are listed with a negative magnitude; in these cases the absolute value of the magnitude was computed from the absolute value of the flux. The 1σ limits of the images are 27.0 (z'), 27.6 (I), 29.2 (R), 28.5 (V), and 29.0 (B).

4. NEAR-INFRARED OBSERVATIONS AND REDUCTION

We have expanded the wide-field HK' coverage of the HDF-N region presented in Barger et al. (1999) with new HK' observations obtained with the University of Hawaii 2.2 m telescope. The new data were obtained UT 2001 March 1–6 with the University of Hawaii QUIRC Camera (QUIRC; Hodapp et al. 1996). QUIRC utilizes a 1024×1024 pixel HgCdTe Astronomical Wide Area Infrared Imaging (HAWAII) array produced by the Rockwell International Science Center. At the f/10 focal ratio the field-of-view of QUIRC is $193'' \times 193''$ with a scale of 0.1886 arcsec pixel $^{-1}$. The observations were made using a notched HK' filter with a central wavelength of $1.8\mu\text{m}$. The filter covers the longer wavelength region of the H -band and the shorter wavelength region of the K -band (roughly the K' filter). Because of its broad bandpass and low sky background, this filter is extremely fast and is roughly twice as sensitive as the H , K' , or K_s filters. Barger et al. (1999) found the empirical relation $HK' - K = 0.13 + 0.05(I - K)$ to convert between the HK' and K bands; for galaxies at most redshifts, $HK' - K \approx 0.3$.

We used a dither pattern of thirteen spatially shifted short exposures of step-size $5'' - 20''$ in all directions. This observing strategy enabled us to use the data images for the subsequent sky subtraction and flat-fielding procedures. Each object is sampled by a different array pixel on each exposure, and the signal-to-noise ratio should improve as the square root of the number of frames. The dark current is insignificant in the QUIRC data. The data were processed using median sky flats generated from the dithered images and calibrated from observations of UKIRT faint standards (Hawarden et al. 2001) taken on the first three nights when the sky was photometric. The FWHM of the total composite image is $0.8''$. The magnitudes were measured in $3.0''$ diameter apertures and corrected to approximate total magnitudes using an offset calculated as the median difference between $3''$ and $6''$ aperture magnitudes measured for a complete $HK' < 20$ sample in the field.

Of the 370 sources in the B01 sample, 276 have HK'

measurements. The corrected aperture HK' magnitudes are given in column 7 of Table 1.




FIG. 3.— (a) R magnitude versus 2–8 keV flux for the CDF-N hard X-ray sample (solid squares; asterisks denote spectroscopically identified stars) and for the Akiyama et al. (2000) data (solid diamonds). The large open squares show the median values of the CDF-N optical magnitudes tabulated in Table 2a. The large open diamond is the median value of the Akiyama et al. magnitudes. (b) R magnitude versus 0.5–2 keV flux for the CDF-N soft X-ray sample (solid squares; asterisks denote spectroscopically identified stars) and for the Lehmann et al. (2001) data (solid diamonds). The large open squares show the median values of the CDF-N optical magnitudes tabulated in Table 2b. The two large open diamonds are the median values of the Lehmann et al. magnitudes. In both (a) and (b) the $\log(f_X/f_R) = 0.0$ (solid) and $\log(f_X/f_R) = \pm 1$ (dashed) lines are loci of constant optical to X-ray flux that cover the range of most of the X-ray selected sources at the brighter X-ray fluxes. Sources fainter than $R = 27.7$ (the 4σ limit) are plotted at this magnitude. Sources saturated in R are shown at a nominal magnitude of $R = 18$ with downward pointing arrows.

5. OPTICAL PROPERTIES OF THE X-RAY SOURCES

In Fig. 3a we plot R magnitude versus 2–8 keV flux for the full CDF-N hard X-ray sample (solid squares). Spectroscopically identified stars are denoted by asterisks. At bright X-ray fluxes we include the Akiyama et al. (2000) *ASCA* Large Sky Survey data (solid diamonds; the two clusters and one source without an optical identification have been excluded, and the star is off-scale), after converting their 2–10 keV fluxes to 2–8 keV assuming $\Gamma = 1.7$ (the value Akiyama et al. assumed in their paper

for the intrinsic photon index). The flux in the R -band is related to the R magnitude by $\log f_R = -5.5 - 0.4R$ (H01). The $\log(f_X/f_R) = 0$ (solid) and $\log(f_X/f_R) = \pm 1$ (dashed) lines are loci of constant optical to X-ray flux that cover the range of most of the hard X-ray selected sources at the brighter X-ray fluxes.

Median optical magnitudes for the CDF-N hard X-ray sample are summarized in Table 2a and are shown in Fig. 3a as large open squares. The large open diamond shows the median optical magnitude for the Akiyama et al. sample. The horizontal bars show the widths of the flux bins while the vertical bars show the 68% confidence range in the median computed using the number of sources in each bin (Gehrels 1986). At bright X-ray fluxes $\log(f_X/f_R) = 0$ roughly matches the median optical magnitudes, presumably because an unobscured AGN dominates the total light output from each source. At fainter X-ray fluxes the medians deviate above $\log(f_X/f_R) = 0$ as the sources become obscured in the optical due to the presence of dust and gas. At the faintest X-ray fluxes the medians flatten as the optical light from the host galaxy begins to dominate the total light output from each source.




FIG. 4.— $B - R$ color versus (a) 2 – 8 keV flux for the hard X-ray sample with $R < 25$ and versus (b) 0.5 – 2 keV flux for the soft X-ray sample with $R < 25$. Both exclude any sources that are saturated. Symbols are as in Fig. 3. (The Lehmann et al. 2001 sample did not include $B - R$ colors so cannot be plotted in b.)

In Fig. 3b we plot R versus 0.5 – 2 keV flux for the full CDF-N soft X-ray sample (solid squares). At bright X-ray fluxes we include the Lehmann et al. (2001) *ROSAT* Ultra Deep Survey data (solid diamonds; groups and clusters and one optically unidentified source have been excluded). Spectroscopically identified stars are denoted by asterisks. Again, the $\log(f_X/f_R) = 0$ (solid) and $\log(f_X/f_R) = \pm 1$ (dashed) lines are loci of constant optical to X-ray flux that cover the range of most of the soft X-ray selected sources at the brighter X-ray fluxes.

Median optical magnitudes for the CDF-N soft X-ray sample are summarized in Table 2b and are shown as large open squares in Fig. 3b. The two large open diamonds show the median optical magnitudes for the Lehmann et al. sample. Figure 3b and Table 2b show that at fluxes above 10^{-15} erg cm $^{-2}$ s $^{-1}$ the $\log(f_X/f_R) = 0$ line matches the median optical magnitudes fairly well, again presumably because an unobscured AGN dominates the total light output from each source. Soft X-ray selected AGN samples at $f_X > 8 \times 10^{-14}$ erg s $^{-1}$ cm $^{-2}$ (0.3 – 3.5 keV) from the *Einstein Observatory* Extended Medium Sensitivity Survey (Stocke et al. 1991) also have values of $\log(f_X/f_R)$ in a similar range. Unlike the hard sample, the medians do not begin to deviate above $\log(f_X/f_R)$ at fainter fluxes, presumably because soft-band sources are mainly unobscured. Below 10^{-15} erg cm $^{-2}$ s $^{-1}$ the median optical magnitudes become constant, within the uncertainties, as the optical light from the host galaxy begins to dominate the total light output from each source.

In Fig. 4a (4b) we plot $B - R$ color versus 2 – 8 keV (0.5 – 2 keV) flux for the CDF-N hard (soft) X-ray sample with $R < 25$, excluding any sources that are saturated. Within the uncertainties the sources redden at faint X-ray fluxes, with the effect being most pronounced in the faint hard X-ray sources. This can be explained by the host galaxy beginning to contribute a larger fraction of the total optical light at these fluxes relative to that contributed by the AGN.

6. SUBMILLIMETER OBSERVATIONS AND REDUCTION

SCUBA jiggle map observations at 850 μ m were taken on the 15 m James Clerk Maxwell Telescope⁶ during observing runs in 2000 April and 2001 March and May. We targeted areas in the CDF-N with large concentrations of X-ray sources to maximize our submillimeter coverage of the field at reasonable sensitivities. The data were obtained in good weather conditions (median $\tau(850\mu\text{m})=0.282$) for a total integration time of 146.5 ks. The maps were dithered to prevent any regions of the sky from repeatedly falling on bad bolometers. The chop throw was fixed at a position angle of 90 deg so that the negative beams would appear 45'' on either side east-west of the positive beam. Regular “skydips” (Lightfoot et al. 1998) were obtained to measure the zenith atmospheric opacities, and the 225 GHz sky opacity was monitored at all times to check for sky stability. Pointing checks were performed every hour during the observations on the blazars 0923+392, 0954+685, 1144+402, 1216+487, 1219+285, 1308+326, or 1418+546. The data were calibrated using jiggle maps of the primary cali-

⁶The James Clerk Maxwell Telescope is operated by the Joint Astronomy Centre on behalf of the UK Particle Physics and Astronomy Research Council, the Netherlands Organization for Scientific Research, and the Canadian National Research Council.




FIG. 5.— Unweighted averages of the normalized spectra for eight spectral types: (a) broad emission lines in the ultraviolet, (b) broad emission lines in the optical, (c) Ly α and strong narrow CIV emission, (d) Ly α emission and CIV absorption, (e) strong [NeIII], (f) [OII] emission, (g) low redshift objects with H α emission, and (h) absorption. The composite spectra are marked with the emission and absorption lines used to identify the individual spectra of Fig. 6. Table 3 lists the number of galaxies per spectral type for objects with spectra given in Fig. 6.

TABLE 2A
MEDIAN OPTICAL MAGNITUDES FOR THE 2 – 8 keV CDF-N SAMPLE

2 – 8 keV flux range (10^{-16} erg cm $^{-2}$ s $^{-1}$)	mean 2 – 8 keV flux (10^{-16} erg cm $^{-2}$ s $^{-1}$)	<i>B</i>	<i>V</i>	<i>R</i>	<i>I</i>	<i>z'</i>
2 – 5	3.35	25.4	24.9	24.5	23.8	23.8
5 – 10	7.55	26.2	25.7	24.5	23.7	23.9
10 – 30	18.2	25.5	24.8	24.2	23.3	23.4
30 – 100	51.4	24.7	23.8	23.2	22.2	22.2
100 – 300	152.	23.8	23.0	22.5	21.7	21.7
300 – 1000	502.	21.1	21.4	20.2	19.9	20.1

TABLE 2B
MEDIAN OPTICAL MAGNITUDES FOR THE 0.5 – 2 keV CDF-N SAMPLE

0.5 – 2 keV flux range (10^{-17} erg cm $^{-2}$ s $^{-1}$)	mean 0.5 – 2 keV flux (10^{-17} erg cm $^{-2}$ s $^{-1}$)	<i>B</i>	<i>V</i>	<i>R</i>	<i>I</i>	<i>z'</i>
2 – 5	4.04	24.9	24.3	23.3	22.4	22.7
5 – 10	7.40	25.0	24.1	22.9	22.0	22.4
10 – 30	17.6	25.2	24.3	23.7	23.0	22.8
30 – 100	55.7	25.5	24.8	24.2	23.5	23.7
100 – 300	174.	24.5	23.9	23.5	22.9	22.9
300 – 1000	493.	22.5	22.1	21.7	21.1	21.2

bration source Mars or the secondary calibration sources CRL618, OH231.8+4.2, or 16293-2422. Submillimeter fluxes were measured using beam-weighted extraction routines that include both the positive and negative portions of the beam profile.

The data were reduced using the dedicated SCUBA User Reduction Facility (SURF; Jenness & Lightfoot 1998). Due to the variation in the density of bolometer samples across the maps, there is a rapid increase in the noise levels at the very edges, so the low exposure edges were clipped. The SURF reduction routines arbitrarily normalize all the data maps in a reduction sequence to the central pixel of the first map; thus, the noise levels in a combined image are determined relative to the quality of the central pixel in the first map. In order to determine the absolute noise levels of our maps, we first eliminated the $\geq 3\sigma$ real sources in each field by subtracting an appropriately normalized version of the beam profile. We then iteratively adjusted the noise normalization until the dispersion of the signal-to-noise ratio measured at random positions became ~ 1 .

The new data were combined in the reduction process with the jiggle maps previously obtained by Barger, Cowie, & Richards (2000) which targeted optically faint radio sources. We also used the Hughes et al. (1998) HDF-N proper ultra-deep map, as rereduced by Barger et al. We restrict our catalog to the 93 arcmin 2 area covered by our SCUBA jiggle maps at better than 5 mJy sensitivity. The area sensitive to 2.5 mJy is 65 arcmin 2 , and the area sensitive to 1.25 mJy is 21 arcmin 2 .

We first measured the submillimeter fluxes at the positions of the 20 cm ($> 5\sigma$) sources from Richards (2000) that fell on our SCUBA maps. Any radio source that we detected above the 3σ level at 850 μ m was included in a “detection list”. We next measured all $> 5\sigma$ submillimeter sources that did not have a radio counterpart. We identified two and added them to the list. The $> 5\sigma$ criterion for the non-radio submillimeter sources ensures that the sources are real and that their positions are relatively accurate, while the weaker $> 3\sigma$ selection criterion is appropriate for a pre-selected radio sample.

After compiling the detection list, we compared the coordinates of our X-ray sources with the coordinates of the sources in the detection list. When an X-ray source was found to lie within $3''$ of a source in the detection list, we identified the list source as the counterpart to the X-ray source. No *Chandra* X-ray source was within $10''$ of either of the two $> 5\sigma$ submillimeter sources without radio counterparts, while all of the radio-detected $> 3\sigma$ submillimeter sources identified as counterparts to X-ray sources were within $1.5''$ of the X-ray source positions; thus, the identifications do not depend critically on the choice of search radius. Once the cross-identifications were made, the sources in the detection list were removed from the SCUBA maps since the wide and complex beam patterns of the bright SCUBA sources can produce spurious detections at other positions. Submillimeter fluxes and uncertainties were then measured at all the unassigned X-ray positions using a recursive loop which removed X-ray

sources that were detected in the submillimeter above the 3σ level in a descending order of significance prior to re-measuring the fluxes. This procedure again avoids multiple or spurious detections of a single bright submillimeter source.

The submillimeter fluxes and uncertainties are given in column 4 of Table 1. The interpretation of the submillimeter data is presented in Barger et al. (2001b).

7. RADIO OBSERVATIONS

We use the Richards (2000) source catalog of the very deep 20 cm VLA map centered on the HDF-N. The image covers a $40'$ diameter region with an effective resolution of $1.8''$ and a 5σ completeness limit of $40 \mu\text{Jy}$. The absolute radio positions are known to $0.1'' - 0.2''$ r.m.s. X-ray sources were identified with radio sources if there were a 5σ radio source within $2''$ of the X-ray position. The use of the radio data in black hole studies is discussed in Barger et al. (2001c) and H01.

8. SPECTROSCOPIC OBSERVATIONS AND REDUCTION

The region surrounding the HDF-N has been the subject of a number of spectroscopic campaigns, and many of the optical sources have known spectroscopic redshifts (e.g., Cohen et al. 2000; Dawson et al. 2001). Additional redshifts have been obtained of the X-ray sample itself (H01; A. E. Hornschemeier et al., in preparation) with the Hobby-Eberly Telescope and with the Low Resolution Imaging Spectrograph (LRIS; Oke et al. 1995) on the Keck⁷ 10 m telescopes. We have obtained our own spectra for as many of the X-ray sources as possible so that we can fully characterize the sources using a homogeneous dataset. In our final spectroscopic sample there are only 7 sources for which we do not have our own spectra, all of which are taken from the Cohen et al. (2000) compilation.

Spectra were primarily obtained with LRIS in multi-slit mode. For sources with $I \leq 24.5$ counterparts within a $2''$ radius, we positioned the slits at the optical centers. For the remaining sources we positioned the slits at the X-ray centroid positions. We used $1.4''$ wide slits throughout. Initially we used the 300 lines mm^{-1} grating blazed at $5000 \text{ \AA}$, which gives a wavelength resolution of $\sim 16 \text{ \AA}$ and a wavelength coverage of $\sim 5000 \text{ \AA}$. The wavelength range for each object depends on the exact location of the slit in the mask but is generally between ~ 5000 and $10000 \text{ \AA}$. After the blue side of LRIS was implemented, we used the 400 lines mm^{-1} grating on the red side and the 600 lines mm^{-1} grism on the blue side, split by the 560 dichroic. This gives a slightly higher resolution spectrum ($\sim 12 \text{ \AA}$) and nearly complete wavelength coverage from ~ 3500 to $10000 \text{ \AA}$. The observations were taken with exposure times of 1 to 1.5 hr per slit mask. Each exposure was broken into three subsets with the objects stepped along the slit by $2''$ in each direction. The sky backgrounds were removed using the medians of the images to avoid the difficult and time-consuming problems of flat-fielding LRIS data. Fainter objects were observed a

number of times with the maximum exposure time for an individual source around 6 hours. Details of the spectroscopic reduction procedures can be found in Cowie et al. (1996).

Additional observations of X-ray sources with bright optical counterparts were obtained with the multifiber wide-field HYDRA spectrograph (Barden et al. 1994) on the WIYN⁸ 3.5 m telescope on UT 2001 February 19–20 and UT 2002 February 6. A 316 lines mm^{-1} grating was used to obtain a wavelength range of $4900 - 10200 \text{ \AA}$ with a resolution of approximately $2.6 \text{ \AA pixel}^{-1}$. The red bench camera was used with the $3''$ ‘red’ HYDRA fibers to maximize the sensitivity at longer wavelengths. To obtain a wavelength solution for each fiber, CuAr comparison lamps were observed in each HYDRA configuration. Our HYDRA masks were designed to maximize the number of optically bright sources in each configuration and to minimize the amount of overlap between configurations. Fibers that were unable to be placed on a source were assigned to a random sky location. We observed 5.3 hours on each of two HYDRA configurations in 2001 and 93 mins on a third configuration in 2002. To remove fiber-to-fiber variations, on-source observations were alternated with $\pm 7.5''$ offset ‘sky’ exposures taken with the same exposure time. This effectively reduced our on-source integration time to 2.6 hours for each configuration in 2001 and to 53 mins for the configuration in 2002.




FIG. 7.— R magnitude versus 2 – 8 keV flux for the hard X-ray sources within a $10'$ radius of the approximate X-ray image center. Solid (open) squares denote sources with (without) redshifts. The dashed line shows the approximate R magnitude to which the sources can be spectroscopically identified with deep 10 m telescope observations ($R = 24$). The star is shown as an asterisk. Objects saturated in R are shown at a nominal magnitude of $R = 18$ with downward pointing arrows. Sources fainter than the 4σ limit of $R = 27.7$ are plotted at this magnitude limit.

Reductions were performed with the standard IRAF⁹ task DOHYDRA. To optimize sky subtraction, we performed a two-step process. In the first step, the DOHYDRA routine was used to create an average sky spectrum

⁷The W. M. Keck Observatory is operated as a scientific partnership among the California Institute of Technology, the University of California, and NASA, and was made possible by the generous financial support of the W. M. Keck Foundation.

⁸The WIYN Observatory is a joint facility of the University of Wisconsin, Indiana University, Yale University, and the National Optical Astronomy Observatory.

⁹IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

TABLE 3
NUMBER OF X-RAY SOURCES PER SPECTRAL TYPE FOR OBJECTS WITH SPECTRA IN FIG. 6

Class	Number
Stars	12
Broad-line emission (ultraviolet or optical)	30
Ly α + CIV emission	9
Ly α emission + CIV absorption	2
[NeIII] emission	22
[OII] or H α emission	67
Absorption	33

using the fibers assigned to random sky locations. This average sky spectrum was then removed from all of the remaining fibers. In the offset images, this step effectively removed all of the sky signal leaving behind only residuals caused by differences among the fibers. To remove these variations, we then subtracted the residuals present in the sky-subtracted offsets from the sky-subtracted on-source spectra. We found this method to be very effective at removing the residuals created by fiber-to-fiber variations with HYDRA.

9. SPECTROSCOPIC REDSHIFTS

Only spectra which could be confidently identified based on multiple emission and/or absorption lines were included in the sample. In Fig. 5 we show unweighted averages of our normalized spectra to illustrate the lines used to identify the different spectral types. Given the positional uncertainties discussed in § 3, we have only cross-identified X-ray sources with spectroscopic counterparts if the radial offsets are $\leq 2''$ (see columns 13 and 14 of Table 1). These redshifts are given in the lower left corner of each thumbnail image in Fig. 2 and are summarized in Table 1. Objects where spectra were obtained but no identification could be made are marked ‘obs’ for observed in column 15 of Table 1. In 13 cases we have spectroscopically identified an object that lies outside our chosen cross-identification radius but still within $5''$ of the X-ray position. We use fainter text in Fig. 2 to give the redshifts for these nearby sources for interest; however, apart from two special cases, we do not give these redshifts in Table 1 nor do we use them in our subsequent analysis. The two special cases are X-ray sources 218 and 290, which appear to lie within the outer envelopes of large extended galaxies with $R_{iso} < 19$. Since there is very little probability that such bright galaxies would randomly fall so close to an X-ray source, we include the redshifts for these two sources in Table 1 and use them in our subsequent analysis.

With this selection 182 of the 370 X-ray sources (49%) have spectroscopic identifications, including 12 stars. The 175 identifiable spectra from the present work (154 galaxies and 6 stars from Keck, 9 galaxies and 6 stars from WIYN) are shown in Fig. 6. (Redshift and B01 catalog number are given at the base of each spectrum. The WIYN spectra are denoted by the letter ‘W’ and are not flux calibrated.) The remaining 7 redshifts are taken from the

compilation of Cohen et al. (2000). In the last column of Table 1 we denote with the letter ‘B’ sources with broad emission lines in their spectra (see Figs. 5a, b). There are 30 such sources in the sample. Table 3 lists the number of galaxies per spectral type (see Fig. 5) for objects with spectra given in Fig. 6.

Outside a $10'$ radius from the approximate *Chandra* image center the X-ray point spread function degrades rapidly, so we do not include these sources in our subsequent figures or analysis. Within $10'$, which corresponds to a 294 arcmin^2 area on the X-ray image, there are 330 *Chandra* sources, including 10 stars. *Hereafter, we refer to this as our X-ray sample.* Of the 189 galaxy sources with $R \leq 24$, 147 (78%) have been spectroscopically identified. The great majority of the identified extragalactic sources with $R \leq 24$ lie below a redshift of one (104/147 or 71%). In Fig. 7 we illustrate for the hard X-ray sample how most sources with $R \leq 24$ can be spectroscopically identified while only a few can be identified at fainter optical magnitudes.




FIG. 8.— Redshift versus R magnitude for the X-ray sources, excluding stars, within a $10'$ radius of the approximate X-ray image center. Objects saturated in R are shown at a nominal magnitude of 18.2. Objects with spectra in Fig. 6 are denoted by solid squares, and the 7 with redshifts from the literature are denoted by open squares. Objects with broad lines are enclosed in a second larger symbol. Objects without redshift identifications are shown at either $z = -0.3$ (open triangles for objects within a $6.5'$ radius of the approximate X-ray image center) or $z = -0.7$ (open diamonds for objects between radii of $6.5'$ and $10'$).

In Fig. 8 we show the redshift-magnitude relation for the galaxy population (squares; unidentified objects are plotted below $z = 0$ with different symbols). Broad-line sources (distinguished by a second larger symbol) are systematically the most optically luminous of the X-ray sources because of the AGN contribution to the light. The gap between $z \sim 1.5$ and 2 reflects the difficulty of identifying sources with redshifts in this range, where [OII] 3727 Å has moved out of the optical window and Ly α 1216 Å has not yet entered.

All of the sources with $z > 1.6$ are either broad-line AGN or have narrow Ly α and/or CIII] 1909 Å emission. There are 14 broad-line AGN versus 11 narrow-line emitters at these redshifts, but this may reflect selection bias in the spectroscopic identifications since the broad-line AGN are generally brighter and easier to identify. Most of the narrow-line sources also have CIV 1550 Å in emission, but two have CIV 1550 Å in absorption.




FIG. 9.— Redshift versus R magnitude for optically selected sources in a $6' \times 6'$ region centered on the HDF-N (open squares). Sources with a counterpart in the X-ray sample are denoted by solid diamonds, and broad-line AGN are enclosed with a second larger symbol.

Intriguingly, while there are a substantial number of optically identified absorption-line sources with $z > 1.6$ known in the HDF-N (e.g., see Brandt et al. 2001b and references therein) and flanking fields, none of these are present in the X-ray sample. We illustrate this in Fig. 9 where we show the redshift-magnitude diagram for an optically selected sample in a $6' \times 6'$ region surrounding the HDF-N. In this region 90% of the 435 sources with $R \leq 23.5$ have been spectroscopically identified (open squares). The sources with a counterpart in the X-ray sample are denoted by solid diamonds. All three of the high-redshift sources with X-ray emission have Ly α in emission (two of these are broad-line AGN), which suggests that only this type of object has X-ray counterparts at our current X-ray sensitivity. Stacking analyses may show a weaker signal below the sensitivity of the individual detections (Brandt et al. 2001b; Nandra et al. 2002).

Roughly a third of the low-redshift sources are detected at 20 cm, and 3 out of the 11 high-redshift narrow emission line sources are detected. However, none of the high-redshift broad-line AGN are detected at 20 cm.




FIG. 10.— (a) Redshift distribution for two bin sizes ($\Delta z = 0.1$ and $\Delta z = 0.01$) for sources within a $10'$ radius of the approximate X-ray image center. (b) R magnitude versus redshift blow-up shows the two structures centered at $z = 0.843$ and $z = 1.0175$. Sources within 1000 km s^{-1} of the center positions are denoted by solid squares, other X-ray sources are denoted by open squares, and field galaxies are denoted by plus signs.

In Fig. 10a we show the histograms of the redshift distribution of our X-ray sample for two bin sizes ($\Delta z = 0.1$ and $\Delta z = 0.01$). The lower resolution binning suggests an excess of objects in two broad redshift bins, which is also evident in the higher resolution binning. In Fig. 10b we show an R magnitude versus redshift blow-up of the two structures centered on $z = 0.843$ and $z = 1.0175$ that are seen in the higher resolution histogram. Sources within 1000 km s^{-1} of the center positions are denoted by solid squares, other X-ray sources are denoted by open squares, and field galaxies, for comparison, are denoted by plus signs. These are the only two structures which contain at least 10 X-ray sources with the specified velocity width. A smaller structure at $z = 0.477$ contains 8 X-ray sources. Figure 10 suggests that there may be large scale structure in the field (spatially the sources also look clustered), but the structures do not dominate the number of X-ray sources in the sample so our redshift distribution should not be strongly affected by clustering. However, the number of sources in these structures (a total of 24 identified sources in the $z = 0.843$ and $z = 1.0175$ structures) is sufficiently large that it could account for a part of the field-

to-field variation seen in the X-ray number counts (e.g., Cowie et al. 2002). Similar redshift structures have also been detected in the CDF-S exposure (Hasinger 2002).

We note that Cohen et al. (2000) also detected strong redshift peaks at $z=0.848$ and $z=1.016$ in their optical survey of the HDF-N region. Dawson et al. (2002) even claimed the serendipitous discovery of ClG 1236+6215, a galaxy cluster with redshift $z=0.85$; however, Bauer et al. (2002) did not significantly detect diffuse X-ray emission associated with this cluster.

10. COLORS AND PHOTOMETRIC REDSHIFTS

Broad-band galaxy colors have been extensively used in recent years to obtain photometric redshift estimates of galaxies (e.g., Bolzonella, Pello, & Miralles 2000). However, the spectral energy distributions of X-ray sources may be rather complex since they arise from both the host galaxy and the AGN. Moreover, the spectral shape of the AGN relative to that of the host galaxy may vary depending on the degree of obscuration. In principle it might be possible to model this with complex templates, but here we simply try to determine empirically which of the sources in our X-ray sample can have their redshifts estimated from the color information using standard galaxy templates. It is precisely for the cases where the AGN is obscured (and hence a spectroscopic redshift hard to obtain) that the photometric redshifts are of most interest, and it is in these cases that the photometric redshift estimates may be expected to work best.

We begin by comparing color-color diagrams with the tracks with redshift that would be followed by a given galaxy type. In Fig. 11a we show $B - R$ versus $R - I$ for the 269 sources in our X-ray sample with $I < 25$, and in Fig. 11b we show $B - I$ versus $I - HK'$ for the 208 sources with $HK' < 21.5$. (In both cases we have excluded sources saturated in the optical images or contaminated by a nearby bright galaxy.) In Fig. 11a circles (squares) denote galaxies with $z < 0.65$ ($z > 0.65$); in Fig. 11b, circles (squares) denote $z < 1$ ($z > 1$). The redshifts $z = 0.65$ and $z = 1$ correspond to where the 4000 Å break passes through the R and I -bands, respectively. We have distinguished sources with broad emission lines in their observed spectra (which may be expected to be dominated by the AGN light) by enclosing them in a second larger symbol. Spectroscopically unidentified galaxies are shown as plus signs. We overlay on the plots the $z = 0$ to $z = 3$ tracks that would be followed by unevolving (i.e., invariant rest-frame spectra) E (top), Sb (middle), and Irr (bottom) galaxies with spectral shapes taken from Coleman, Wu, & Weedman (1980).

From Fig. 11a we see that nearly the whole X-ray sample lies either in the region populated by the galaxy tracks or in regions where a dominant AGN would migrate the colors. The broad-line systems are all extremely blue and lie in a small region of the $B - R$ versus $R - I$ plane. However, they substantially overlap with the irregular galaxies, making it hard to distinguish between low redshift irregulars and luminous high-redshift AGN purely on the basis of color. The redder galaxies fall in two well-defined regions. The reason for the split is the passage of the 4000 Å break through the R -band at this redshift, which results in the colors, which were originally moving up the solid arms of

the unevolving galaxy tracks, crossing to the dashed arms and then migrating down with redshift. There appear to be relatively few sources with colors as red as an elliptical galaxy at $z > 0.65$ but many sources with colors similar to Sb or Sa galaxies. There are substantial numbers of spectroscopically unidentified galaxies in both of the well-defined regions and also in the AGN/Irr region, but there are only a tiny number outside those areas.




FIG. 11.— (a) $B - R$ versus $R - I$ for the $I < 25$ X-ray sources within a $10'$ radius of the approximate X-ray image center, excluding the sources that are saturated or contaminated by a bright neighbor. Plus signs denote galaxies without redshift identifications. Circles (squares) denote galaxies with $z < 0.65$ ($z > 0.65$). Broad-line sources are enclosed in a second larger symbol. Objects with $R - I > 1.9$ ($B - R > 3.4$) are plotted at $R - I = 1.9$ ($B - R = 3.4$). The overlaid curves show the tracks that would be followed by unevolving E (top), Sb (middle), and Irr (bottom) galaxies with spectral shapes from Coleman, Wu, & Weedman (1980). The redshift range of the tracks is $z = 0$ to $z = 3$, with the $z < 0.65$ ($z > 0.65$) region illustrated by a solid (dashed) line. (b) $B - I$ versus $I - HK'$ for the $HK' < 21.5$ X-ray sources within the $10'$ radius, excluding the sources that are saturated or contaminated by a bright neighbor. The symbols and the galaxy tracks are the same as in (a), except here the circles (squares) denote galaxies with $z < 1$ ($z > 1$), and the solid (dashed) tracks illustrate the region with $z < 1$ ($z > 1$). Objects with $I - HK' > 5.9$ ($B - I > 5.9$) are plotted at $I - HK' = 5.9$ ($B - I = 5.9$). The dotted line shows the star-galaxy discriminator of Gardner (1992), modified to the HK' system (Barger et al. 1999). Stars are expected to lie to the left of this line, as is the case for the one spectroscopically identified star which is not saturated (asterisk).

The unidentified sources in Fig. 11b seem to preferen-

tially lie at $z > 1$, with only a small number at lower redshifts. This is partially a selection effect since the higher redshift galaxies have redder NIR colors and hence are more likely to show up in an HK' magnitude-limited sample. As in Fig. 11a, very few of the galaxies lie near the elliptical track at high redshifts, with most of the sources having Sa/Sb colors or bluer. There is a substantial spread in the $I - HK'$ colors of the broad-line sources, which may reflect varying host galaxy contributions in the NIR. In Fig. 11b we show the star-galaxy discriminator (dotted line) used by Gardner (1992) and modified to the HK' system (Barger et al. 1999). Stars are expected to lie to the left of this line, and it is clear from the figure that there are very few optically faint stars in the X-ray sample (most have $I \lesssim 19$), with only one (spectroscopically identified) star lying in this region (asterisk).

We next used the Hyperz code of Bolzonella et al. (2000) and the magnitudes (including HK' , where available) of Table 1 to estimate photometric redshifts. Seven updated Bruzual & Charlot (1993) models corresponding to present-day E, S0, Sa, Sb, Sc, Sd, and Irr galaxies were used, as well as a single-burst model. These templates cover the entire range of possible ages. The ratio of the photometric redshift estimate from Hyperz to the observed spectroscopic redshift is shown versus $B - I$ color in Fig. 12 for spectroscopically identified sources which are not saturated in any of the optical bands or contaminated by neighbors. Broad-line sources are enclosed in a second larger symbol. As we saw from Fig. 11, the present analysis cannot distinguish the AGN from the Irr galaxies. However, redder than $B - I = 1.5$ (vertical dotted line), the photometric redshift estimates become more robust, and most sources have photometric redshifts within 25% of the spectroscopic redshift, as shown by the horizontal dashed lines.




FIG. 12.— Ratio of photometric (calculated with the Hyperz program and the magnitudes in Table 1) to spectroscopic redshifts versus $B - I$ color for the sources within a $10'$ radius of the approximate X-ray image center, excluding the sources that are saturated or contaminated by a bright neighbor. Broad-line sources are enclosed with a second larger symbol. AGN contamination degrades the photometric redshift fits, but redder than $B - I = 1.5$ (vertical dotted line) the photometric redshifts generally agree with the actual redshifts to within 25% (horizontal dashed lines).

The use of compactness and apparent magnitude as prior constraints can improve the agreement between the

photometric and spectroscopic redshifts for sources with blue colors since AGN are likely to be at high redshifts, compact, and bright, while the Irr galaxies are likely to be at low redshifts, extended, and faint. However, even with these constraints, the featurelessness of the blue spectra and the overlap of the apparent magnitudes means there is still a high degree of indeterminacy for the blue objects. Ultraviolet data would significantly improve the identification of these types of objects.

In our subsequent analysis we restrict our use of the photometric redshifts to galaxies with $B - I > 1.5$ and $I < 25$ that are also not saturated or near a contaminating bright object.




FIG. 13.— Spectroscopic redshift versus I magnitude (solid squares) for the (a) 2 – 8 keV and (b) 0.5 – 2 keV X-ray samples within a $10'$ radius of the approximate X-ray image center. Broad-line sources are enclosed in a second larger symbol. The open symbols denote photometric redshifts for unidentified or unobserved sources with $B - I > 1.5$. In (a) 22 objects bluer than $B - I = 1.5$ without spectroscopic redshifts are omitted from the plot and in (b) 26. The solid lines show the tracks of unevolving Sb galaxies with absolute rest-frame magnitudes $M_I = -22$ (lower tracks) and $M_I = -24.5$ (upper tracks).

11. ABSOLUTE MAGNITUDES OF THE OPTICAL COUNTERPARTS

In Figs. 13 we show redshift versus I magnitude for our 0.5 – 2 keV and 2 – 8 keV samples; we plot only objects having optical magnitudes between $I = 20$ and $I = 25$. Sources with spectroscopic (photometric) red-

shifts are denoted by solid (open) symbols. Broad-line sources are enclosed in a second larger symbol. The solid lines show the expected tracks for an unevolving Sb galaxy with $M_I = -24.5$ (upper tracks) and $M_I = -22$ (lower tracks). The $M_B^* = -20.4$ galaxy from Loveday et al. (1992) roughly corresponds to an Sb galaxy with $M_I^* = -22.0$.

Of the 236 hard (2–8 keV) X-ray sources in our sample, 177 lie in the $I = 20$ to $I = 25$ range, of which 155 (97 spectroscopic and 58 photometric) are shown (22 $B - I < 1.5$ objects are omitted). Of these, 29 are brighter than $M_I = -24.5$ and 113 are brighter than $M_I^* = -22$. Of the 288 soft X-ray sources in our sample, 205 lie in the $I = 20$ to $I = 25$ range, of which 179 (115 spectroscopic and 64 photometric) are shown (26 $B - I < 1.5$ objects are omitted). Of these, 34 are brighter than $M_I = -24.5$ and 134 are brighter than $M_I^* = -22$. Thus, most of the galaxies in both the hard and soft samples are comparable to or more luminous than M_I^* .

12. X-RAY PROPERTIES

In our analysis of the X-ray source properties, we consider two restricted uniform flux-limited subsamples. *Hereafter, we refer to these as our ‘deep’ and ‘bright’ subsamples.* For the deep subsample we consider a 6.5′ radius high image quality and high exposure time region around the approximate center of the X-ray image. The minimum exposure time for a source in the 6.5′ radius region is 660 ks. The 90% enclosed energy radius is less than 7 arcseconds for the 2–8 keV band. Allowing for these variations, we select sources detected above fluxes of 10^{-16} erg cm $^{-2}$ s $^{-1}$ (0.5–2 keV), 5×10^{-16} erg cm $^{-2}$ s $^{-1}$ (2–8 keV), and 10^{-15} erg cm $^{-2}$ s $^{-1}$ (4–8 keV). For the bright subsample we consider the 10′ radius region, except now we select only sources detected above fluxes of 10^{-15} erg cm $^{-2}$ s $^{-1}$ (0.5–2 keV), 5×10^{-15} erg cm $^{-2}$ s $^{-1}$ (2–8 keV), and 10^{-14} erg cm $^{-2}$ s $^{-1}$ (4–8 keV). The incompleteness and bias corrections are expected to be small at the flux levels of both the deep and bright subsamples (Cowie et al. 2002).

12.1. Redshift Distributions

We show the redshift distributions for the deep (squares) and bright (inverted triangles) subsamples versus X-ray flux in Figs. 14. Broad-line sources are enclosed in a second larger symbol. The spectroscopically unidentified or unobserved sources with photometric redshift estimates from § 10 are plotted as open symbols at those redshifts, while sources without a photometric redshift estimate are plotted below the $z = 0$ line. In Fig. 14a we also include the *ASCA* Large Sky Survey data (circles) from Akiyama et al. (2000) and the SSA13 (diamonds) and A370 (triangles) data from Barger et al. (2001a, b). In Fig. 14b we include the *ROSAT* Ultra Deep Survey data (circles) from Lehmann et al. (2001).

The percentage of the total light in the hard or soft deep subsample that comes from spectroscopically identified point sources in each redshift interval is indicated. In the hard deep subsample at least 35% arises below $z = 1$ and at least 55% arises below $z = 2$. These percentages increase to 40% and 73% if we include the sources in the hard deep subsample with photometric redshifts. Similarly, in the soft deep subsample at least 31% arises below

$z = 1$ and at least 62% arises below $z = 2$. These percentages increase to 36% and 73% if we include the sources in the soft deep subsample with photometric redshifts. Thus, the bulk of the total 2–8 keV and 0.5–2 keV flux in the deep subsamples arises at recent times, which is broadly consistent with the results of other X-ray samples (Barger et al. 2001c; Hasinger 2002).




FIG. 14.— Redshift versus X-ray flux for the deep (squares) plus bright (inverted triangles) (a) hard and (b) soft X-ray subsamples (see § 12). Broad-line sources are enclosed in a second larger symbol. Spectroscopically unidentified or unobserved sources with (without) photometric redshifts are plotted as open symbols at those redshifts (below the $z = 0$ line). In (a) we also plot the Akiyama et al. (2000) data (circles) and the Barger et al. (2001a,b) SSA13 (diamonds) and A370 data (triangles). In (b) we include the Lehmann et al. (2001) data (circles). At the left is given the percentage of the total light in the hard or soft deep subsample that comes from spectroscopically identified point sources in each redshift interval.

12.2. Spectral Characteristics

The conversion from counts to flux, the correction for opacity effects, and the K -correction all depend on the shape of the source spectrum. For a power-law spectrum with photon index Γ the counts s $^{-1}$ in an energy band E_1 to E_2 are given by $N_{E_1-E_2} = \kappa \int_{E_1}^{E_2} A(E) E^{-\Gamma} dE$, where $A(E)$ is the effective detector area at energy E and $\kappa E^{-\Gamma}$ has units of photons cm $^{-2}$ s $^{-1}$ keV $^{-1}$. Once the normalization of the spectrum, κ , is determined from the observed

counts, the flux is $f = \int_{E_1}^{E_2} F(E) dE$ where $F = \kappa E^{1-\Gamma}$. The energy index is $\alpha = \Gamma - 1$.

For their counts to flux conversion, B01 used an effective Γ for each source determined from the ratio of the hard-band to soft-band counts. In this section we take a different approach and try to determine the best choice of input spectrum by considering the 4–8 keV sample, which is the least biased by any optical depth selection.

We consider only sources within $6.5'$ of the approximate X-ray image center that have 4–8 keV fluxes above 2×10^{-15} erg cm $^{-2}$ s $^{-1}$. All the sources that satisfy these criteria are also detected in the 0.5–2 keV and 2–8 keV bands. Moreover, since the selected flux limit is well below the value 10^{-14} erg cm $^{-2}$ s $^{-1}$ where most of the XRB contributions originate (e.g., Cowie et al. 2002), these sources should be representative of the ‘typical’ source dominating the background.




FIG. 15.— Observed $(0.5 - 2 \text{ keV counts})/(4 - 8 \text{ keV counts})$ versus $(2 - 4 \text{ keV counts})/(4 - 8 \text{ keV counts})$. The solid squares show a 4–8 keV sample within $6.5'$ of the approximate X-ray image center with fluxes greater than 2×10^{-15} erg cm $^{-2}$ s $^{-1}$. All of the sources in this sample are also detected in the 0.5–2 keV and 2–8 keV bands. The large open squares (diamonds) show sources with spectroscopic redshifts $z < 1$ ($z > 1$). The solid line is the track for an unobscured power-law source with varying photon index; the tick marks correspond to $\Gamma = (-1, -0.5, 0, 0.5, 1, 1.5, 2, 2.5)$ with the values increasing from bottom to top. The dashed line shows the track of a power-law source with an intrinsic $\Gamma = 2$ versus rest-frame N_H ; the tick marks correspond to $\log N_H = (21, 21.5, 21.7, 22, 22.5, 22.7)$ with the values increasing from top to bottom. The two dotted lines show the tracks of similar absorbed power-laws with intrinsic spectral photon indices $\Gamma = 1.5$ (left) and $\Gamma = 2.5$ (right).

In Fig. 15 we show $(0.5 - 2 \text{ keV counts})/(4 - 8 \text{ keV counts})$ versus $(2 - 4 \text{ keV counts})/(4 - 8 \text{ keV counts})$. (The 2–4 keV counts were determined from the difference of the 2–8 keV and 4–8 keV counts.) Sources with spectroscopic redshifts are denoted by a second larger symbol (squares for $z < 1$ and diamonds for $z > 1$). Two simple models are also shown. The solid line is the track expected for a power-law source with varying photon index Γ . A single photon index cannot simultaneously match both the hard and soft X-ray colors, and a strong break from harder to softer spectra is required as the energy increases. An only slightly more complicated model is one where we assume a fixed intrinsic photon index that is absorbed at low en-

ergies by photoelectric absorption from a varying column density of material in front of the source. The dashed line is the track expected for a power-law source with a fixed $\Gamma = 2$ index and a varying opacity, computed using the cross-section for solar abundances (Morrison & McCammon 1983). Over the energy range of interest for significant opacity, the cross-section, $\sigma(E)$, can be approximated by a single power-law

$$\sigma(E) = 2.4 \times 10^{-22} E^{-2.6} \text{ cm}^2$$

with E in keV. The dotted lines show similar absorbed power-laws with intrinsic photon indices $\Gamma = 1.5$ (left line) and $\Gamma = 2.5$ (right line). As a consequence of the power-law energy dependence of the cross-section, the effective hydrogen column density has a redshift dependence

$$N_{eff} = N_H / (1 + z)^{2.6}$$

It is clear from Fig. 15 that the X-ray colors of the sources are reasonably well-described by this type of model and that most of the sources lie within the $\Gamma = 1.5$ to $\Gamma = 2.5$ tracks. The fact that the dashed line in Fig. 15 is nearly vertical shows that absorption effects have little consequence above 2 keV but produce significant flux suppressions below 2 keV.




FIG. 16.— Histograms for the bright and deep subsamples selected in the 2–8 keV band showing the distributions by optical spectral type (see Table 3) of the hardness of the sources, as specified by the $(0.5 - 2 \text{ keV})/(2 - 8 \text{ keV})$ counts ratio. The counts ratio for any source not detected in the soft band was determined using the 0.5–2 keV upper limit given in Table 3 of B01, so such sources could lie to the left of the current positions. In the third panel from the bottom the solid (dashed) line shows the distribution of sources with [NeIII] emission (CIV emission).

In Fig. 16 we show for the bright and deep subsamples selected in the 2–8 keV band the hardness distribution (as specified by the ratio of the soft-band to hard-band counts) with optical spectral type. The broad-line sources have an extremely narrow hardness range which would correspond to a spectral index of $\Gamma = 1.7$ in the absence of optical depth effects. Nearly all have $\Gamma < 1.8$. The stars are the softest X-ray sources, as is well-known. The remaining sources span a wide range of hardness, from very hard to comparable to that of the broad-line AGN. The line-of-sight column densities for these latter sources are only loosely related to the optical spectral characteristics;

this is likely due to different viewing angles for AGN leading to different X-ray and optical obscurations. Based on the distribution of the counts ratios of the broad-line AGN in Fig. 16, we adopt an intrinsic $\Gamma = 1.8$ in the absence of absorption in the subsequent discussion.

We now use the $\Gamma = 1.8$ input spectrum to investigate the range of intrinsic column densities N_H (i.e., the column densities in the rest-frame of each source) that produce the absorption. In Fig. 17 we plot $(0.5 - 2 \text{ keV counts})/(2 - 8 \text{ keV counts})$ versus redshift for the $0.5 - 2 \text{ keV}$ and $2 - 8 \text{ keV}$ deep subsamples. Sources with spectroscopic (photometric) redshifts are denoted by solid (open) symbols. We overlay on the data fixed intrinsic N_H curves generated assuming an intrinsic $\Gamma = 1.8$ power-law spectrum and absorption by the photoelectric cross-section. We do not show any curves after Compton thickness effects become important. At $z < 1$ the sources in the sample appear to be well-described by the $\log N_H \leq 23.5$ curves. At $z > 1$ there are only a small number of spectroscopically unidentified sources which potentially could have column densities much in excess of $\log N_H = 23.5$. Such sources may be very optically faint and thus preferentially lack redshifts.




FIG. 17.— The natural logarithm of the $(0.5 - 2 \text{ keV})/(2 - 8 \text{ keV})$ counts ratio versus redshift for the $0.5 - 2 \text{ keV}$ and $2 - 8 \text{ keV}$ deep subsamples. Sources only detected in the soft (hard) band are shown with upward (downward) pointing arrows using the limits from Table 3 of B01. We overlay as solid curves the values expected for a source with an intrinsic $\Gamma = 1.8$ spectrum (shown by the dashed line) and an intrinsic column density of gas and dust corresponding to $\log N_H = (22, 22.5, 23, 23.5)$ (solid lines). Sources with spectroscopic redshifts are denoted by solid symbols. Unidentified or unobserved sources are denoted by open symbols at either the photometric redshift or at $z = 1.5$, if the object is too faint or too blue to have a photometric redshift.

In Fig. 18 we plot the deduced intrinsic N_H values for the individual sources in the $2 - 8 \text{ keV}$ selected bright and deep subsamples with either spectroscopic or photometric redshifts. The inferred N_H values range from about 10^{21} cm^{-2} to $5 \times 10^{23} \text{ cm}^{-2}$. As expected, the broad-line AGN (denoted by a second larger symbol) have comparatively low N_H values.

Our description is simple, and we caution that when sources get absorbed by material that is almost or fully Compton-thick, other components could take over to determine the spectral shape, including scattering of X-rays around the absorber, multiple lines-of-sight through the

absorber, circumnuclear starburst emission, etc. These effects will tend to give more flux at low energies than would be expected from the simple model. Thus, column densities can be underestimated, with the underestimation factor getting larger as the column density gets larger. Consequently, there could still be some Compton-thick sources in our sample, even if we do not recognize them as such. This is an interesting issue, and in the next section we examine whether the XRB *requires* contributions from a population of Compton-thick sources.




FIG. 18.— Rest-frame column densities required to produce the observed $(2 - 8 \text{ keV})/(0.5 - 2 \text{ keV})$ counts ratios versus redshift for the $2 - 8 \text{ keV}$ bright and deep subsamples with spectroscopic (solid) or photometric (open) redshifts. Sources not detected in the soft band are shown with upward pointing arrows using the limits from Table 3 of B01. The $z = 5.18$ source is off-scale. Sources with broad emission lines in their spectra are indicated with a second larger symbol. These broad-line sources are among the lowest column density systems.

12.3. High Energy XRB Contributions

An important question is how much of the very high energy ($> 8 \text{ keV}$) XRB our current sample has resolved, and hence how much light might still be needed from Compton-thick sources. In Fig. 19 we plot the total XRB contributions from the $2 - 8 \text{ keV}$ deep subsample (thick solid line), computed for each source assuming an intrinsic $\Gamma = 1.8$ power-law together with an optical depth determined from the $(0.5 - 2 \text{ keV})/(2 - 8 \text{ keV})$ counts ratio. The dashed (Boldt 1987) and dotted (Gruber et al. 1984; Gruber 1992; Fabian & Barcons 1992) lines show the XRB measurements obtained by *HEAO-A*. The deep subsample produces about 87% of this background at 3 keV and 57% at 20 keV .

However, if the *Chandra* data were renormalized at 3 keV to the higher *ASCA* or *BeppoSAX* measurements (the latter, taken from Vecchi et al. 1999, are shown in Fig. 19 by the dot-dashed line), then the shape of the combined contributions from the *Chandra* sources matches fairly well that of the observed XRB (thin solid line) at both energies. Thus, whether the match to the high energy XRB requires either a substantial population of as-yet undetected Compton-thick sources or some change in the spectral shape of the current sources from the simple power-law dependence clearly depends critically on how the low energy and high energy XRB measurements tie

together (see also Fabian, Wilman, & Crawford 2002 for a theoretical discussion).




FIG. 19.— Total XRB contribution from the 2 – 8 keV deep subsample (thick solid line) based on an intrinsic $\Gamma = 1.8$ power-law dependence combined with an opacity determined from the $(0.5 - 2 \text{ keV})/(2 - 8 \text{ keV})$ counts ratio for each source in the subsample. The dashed (dotted) line shows the *HEAO-A* XRB measurement from Boldt 1987 (Gruber et al. 1984; Gruber 1992; Fabian & Barcons 1992). The dot-dashed line is the *BeppoSAX* measurement of Vecchi et al. (1999). Other measurements in the 1 – 10 keV range lie between the *HEAO-A* and *BeppoSAX* measurements. The thin solid line is the total XRB contribution from the 2 – 8 keV deep subsample, renormalized to match the *BeppoSAX* measurement at 3 keV.

12.4. X-ray Luminosities

Since the translation of observed *Chandra* counts to flux depends on the assumed spectral shape, the computation of the intrinsic rest-frame X-ray luminosities using the intrinsic $\Gamma = 1.8$ spectrum and the inferred rest-frame N_H values requires a recomputation of the B01 fluxes, followed by a correction for the opacity (which reduces the observed flux) and a $(1+z)^{-0.2}$ K -correction. It turns out that all of these effects are small in the 2 – 8 keV band, and the final 2 – 8 keV luminosities that we calculate only differ from those which would be computed directly with the B01 fluxes and a zero K -correction by an average factor of 0.83 with a weak redshift dependence. This factor is similar to the 0.85 constant factor used by Barger et al. (2001a) to empirically correct their 2 – 8 keV luminosities for these effects.

In Fig. 20 we show our rest-frame opacity-corrected and K -corrected 2 – 8 keV luminosities versus redshift for the 2 – 8 keV bright and deep subsamples with spectroscopic identifications. Figure 20 shows very little evolution in the maximum X-ray luminosities with decreasing redshifts until $z \lesssim 0.5$, at which point the volume becomes too small to probe effectively very high luminosity sources. If there were considerable evolution of the maximum luminosities, as is seen in the optical, then we would not have detected any high luminosity sources at low redshift.

There has been some discussion in the literature about rare finds of high-redshift type II quasars (e.g., Norman et al. 2002; Stern et al. 2002a). We note that at $L_{2-8 \text{ keV}} \gtrsim 10^{44} \text{ erg s}^{-1}$ and $z > 2$ we also find two such non-broad-line systems, objects 184 and 287 from Table 1, both of which are narrow $\text{Ly}\alpha$ emitters. At $z \approx 1$ we

also see two such systems, objects 69 and 280 in Table 1, both of which have $[\text{NeIII}]$ and $[\text{NeV}]$ emission; 280 also has narrow Balmer emission lines. However, we stress that high-redshift type II quasars are by no means responsible for producing the bulk of the XRB at these energies (see § 12.1).




FIG. 20.— Rest-frame opacity-corrected and K -corrected 2 – 8 keV luminosity versus redshift for the hard deep and bright subsamples with spectroscopic identifications. The solid squares (diamonds) denote sources in the deep (bright) subsample. Sources identified spectroscopically as broad-line AGN are denoted by a second larger symbol. The solid (dashed) curve shows the X-ray luminosity limit inferred from the deep (bright) subsample flux limit of $5 \times 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$ ($5 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$).

Because the optical depth effects are much larger at low energies, the corrections described above depend sensitively on the choice of intrinsic spectrum; thus, we have not attempted to calculate intrinsic rest-frame 0.5 – 2 keV luminosities. Instead, in Fig. 21 we show observed-frame 0.5 – 2 keV luminosity (based on the B01 fluxes with no opacity correction) versus redshift for the soft bright and deep subsamples at $z < 1$. Sources with an intrinsic rest-frame $L_{2-8 \text{ keV}} \gtrsim 10^{42} \text{ erg s}^{-1}$ are denoted by open symbols, and those with fainter hard X-ray luminosities are denoted by solid symbols. X-ray luminosities of local starbursts never exceed $10^{42} \text{ erg s}^{-1}$ (Zezas, Georgantopoulos, & Ward 1998; Moran et al. 1999), so any source more luminous than this is very likely an AGN.

Below $10^{42} \text{ erg s}^{-1}$ the sources could be more ‘typical’ galaxies whose X-ray emission is dominated by a population of X-ray binaries, hot interstellar gas, or a low-luminosity AGN. We detect 18 low-luminosity sources in the 0.5 – 2 keV deep subsample, which corresponds to a surface density of $490^{+140}_{-110} \text{ deg}^{-2}$. These sources contribute a very small fraction (approximately 3%) of the total 0.5 – 2 keV light in the deep subsample and are not a significant contributor to the XRB. A more thorough discussion of the optically bright, X-ray faint source population to which normal galaxies belong can be found in A. E. Hornschemeier et al., in preparation.




FIG. 21.— Observed 0.5–2 keV luminosity versus redshift for the soft deep and bright subsamples with spectroscopic identifications. The open squares (diamonds) denote sources in the deep (bright) subsample with $L_{2-8 \text{ keV}} \gtrsim 10^{42} \text{ erg s}^{-1}$; the solid symbols denote those with fainter hard X-ray luminosities. Sources identified spectroscopically as broad-line AGN are denoted by a second larger symbol. The solid (dashed) curve shows the observed X-ray luminosity limit inferred from the deep (bright) subsample flux limit of $10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$ ($10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$).

12.5. Optical versus X-ray Luminosities

We calculate B -band luminosities, L_B , for the X-ray sources using the relation

$$L_B = 4\pi d_L^2 (1.5 \times 10^{14}) f_{4500(1+z)} (1+z)^{-1}$$

where d_L is the luminosity distance in cm, $f_{4500(1+z)}$ is the rest-frame B -band flux in units of $\text{erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1}$ (estimated by interpolating from the observed fluxes corresponding to the magnitudes given in Table 1), and $1.5 \times 10^{14} \text{ Hz}$ is the width of the B -band filter. In Fig. 22 we plot rest-frame L_B versus $L_{2-8 \text{ keV}}$ (from § 12.4) for the 2–8 keV bright and deep subsamples. The solid lines have slopes $\log(L_B/L_{2-8 \text{ keV}}) = 100, 10, 1, 0.1$, and 0.01 . The open (solid) symbols denote sources with redshifts $z > 1.2$ ($z < 1.2$). The second larger symbols denote spectroscopic broad-line AGN.

The $z > 1.2$ sources approximately follow the line $\log(L_B/L_{2-8 \text{ keV}}) = 1$, which suggests that the central AGN in these galaxies dominate their optical luminosities. At $L_{2-8 \text{ keV}} \lesssim 10^{43} \text{ erg s}^{-1}$ the locus flattens above the $\log(L_B/L_{2-8 \text{ keV}}) = 1$ line; here the host galaxy light starts to dominate the total optical light. At the high X-ray luminosities most of the identified sources are broad-line AGN. Since the unobscured AGN light dominates the optical light in these systems, no matter what redshifts the sources are at we should have no trouble identifying them spectroscopically. However, at high redshifts and high X-ray luminosities, we may be unable to spectroscopically identify high-extinction AGN due to the obscuration of the optical light originating from the AGN.




FIG. 22.— Rest-frame B -band luminosity versus rest-frame opacity-corrected and K -corrected 2–8 keV luminosity for the spectroscopically identified hard deep and bright subsamples. The open (solid) symbols denote sources with redshifts $z > 1.2$ ($z < 1.2$). The second larger symbols denote spectroscopic broad-line AGN. The two objects which are saturated in R and I are shown as lower limits with upward pointing arrows. The solid lines have slopes $\log(L_B/L_{2-8 \text{ keV}}) = 100, 10, 1, 0.1$, and 0.01 . The dotted line is the luminosity of an $M_B^* = -20.4$ galaxy from Loveday et al. (1992).

13. SUMMARY

We presented a catalog of the optical, NIR, submillimeter, and radio properties of the X-ray sources identified in the $\approx 1 \text{ Ms}$ *Chandra* exposure of the CDF-N. We now have redshifts for 182 of the 370 X-ray sources, and we presented the spectra for 175. The redshift identifications are very complete (78%) for the $R \leq 24$ galaxy sources in a $10'$ radius around the approximate X-ray image center. All of the X-ray sources with $z > 1.6$ are either broad-line AGN or have narrow $\text{Ly}\alpha$ and/or $\text{CIII}] 1909 \text{ \AA}$ emission; none of the known $z > 1.6$ absorption line systems in the field are detected in X-rays.

We found spectroscopic evidence for large scale structure in the field which could account for a part of the field-to-field variation seen in the X-ray number counts. However, since the observed structures do not dominate the number of X-ray sources in the sample, the redshift distribution should not be too strongly affected by clustering.

The broad-line sources are all extremely blue, making it hard to distinguish between low redshift irregulars and luminous high-redshift AGN purely on the basis of color. Very few of the redder X-ray sources, whether spectroscopically identified or not, lie outside the regions populated by unevolving spiral galaxy tracks.

We estimated photometric redshifts for the sources using our five broad-band colors (six when HK' was available). Above $B - I = 1.5$ the photometric redshift estimates are quite robust; for most sources with available spectroscopic redshifts the photometric redshifts are within 25%. The majority of galaxies have absolute magnitudes comparable to or more luminous than $M_I^* = -22$.

We found that the spectral energy distributions of the X-ray sources are well-described by a $\Gamma = 1.8$ intrinsic spectrum corrected for absorption. We determined the absorbing column densities for our sources with red-

shifts and found that they range from about 10^{21} cm^{-2} to $5 \times 10^{23} \text{ cm}^{-2}$.

We calculated intrinsic rest-frame hard and observed-frame soft X-ray luminosities. We found very little evolution in the maximum hard X-ray luminosities until $z \lesssim 0.5$, at which point the volume becomes too small to probe the very high luminosity sources. At soft X-ray luminosities of $L_{0.5-2 \text{ keV}} < 10^{42} \text{ erg s}^{-1}$ the ‘typical’ galaxy, whose X-ray emission may be dominated by a population of X-ray binaries, hot interstellar gas, or a low-luminosity AGN, becomes important.

We calculated *B*-band luminosities, L_B , for the X-ray sources. Sources with $z > 1.2$ approximately follow the line $\log(L_B/L_{2-8 \text{ keV}}) = 1$, which suggests that the AGN dominate the optical luminosities. At $L_{2-8 \text{ keV}} \sim 10^{43} \text{ erg s}^{-1}$ the locus flattens above the line, which suggests this is where the host galaxy light starts to dominate the optical light.

Redshift slices of the sources versus X-ray flux show that substantial contributions to the total X-ray flux in both bands (of order one-third using only the present spectroscopic sample) come from $z < 1$. Thus, major accretion onto supermassive black holes has occurred since the Universe was half its present age.

We estimated that the *Chandra* sources that produce

87% of the *HEAO-A* XRB at 3 keV produce 57% at 20 keV, provided that at high energies the spectral shape of the sources continues to be well-described by a $\Gamma = 1.8$ power-law. However, when the *Chandra* contributions are renormalized to the *BeppoSAX* XRB at 3 keV, the shape matches fairly well the observed XRB at both energies. Thus, whether a substantial population of as-yet undetected Compton-thick sources or some change in the spectral shape of the current sources from the simple power-law dependence is required to resolve the high energy XRB depends critically on how the low energy and high energy XRB measurements tie together.

We thank the referee, David Helfand, for helpful comments that improved the manuscript. We thank Andy Fabian, Scott Tremaine, and Omar Almaini for useful discussions. We gratefully acknowledge support from the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation (AJB), NSF grants AST-0084847 (AJB, PI) and AST-0084816 (LLC), NASA grants DF1-2001X (LLC, PI), NAS8-01128 (GPG, PI), and G02-3187A (WNB), NSF CAREER award AST-9983783 (WNB), NASA GSRP grant NGT 5-50247 (AEH), and the Pennsylvania Space Grant Consortium (AEH).

REFERENCES

- Akiyama, M., et al. 2000, *ApJ*, 532, 700
 Alexander, D. M., Brandt, W. N., Hornschemeier, A. E., Garmire, G. P., Schneider, D. P., Bauer, F. E., & Griffiths, R. E. 2001, *AJ*, 122, 2156
 Baldi, A., Molendi, S., Comastri, A., Fiore, F., Matt, G. & Vignali, C. 2002, *ApJ*, 564, 190
 Barden, S. C., Armandroff, T., Muller, G., Rudeen, A. C., Lewis, J., & Groves, L. 1994, *Instrumentation in Astronomy VIII*, ed. D.L. Crawford and E.R. Craine, SPIE Vol. 2198, p. 87.
 Barger, A. J., Cowie, L. L., Trentham, N., Fulton, E., Hu, E. M., Songaila, A., & Hall, D. 1999, *AJ*, 117, 102
 Barger, A. J., Cowie, L. L., & Richards, E. A. 2000, *AJ*, 119, 2092
 Barger, A. J., Cowie, L. L., Mushotzky, R. F., & Richards, E. A. 2001a, *AJ*, 121, 662
 Barger, A. J., Cowie, L. L., Steffen, A. T., Hornschemeier, A. E., Brandt, W. N., & Garmire, G. P. 2001b, *ApJ*, 560, L23
 Barger, A. J., Cowie, L. L., Bautz, M. W., Brandt, W. N., Garmire, G. P., Hornschemeier, A. E., Ivison, R. J., & Owen, F. N. 2001c, *AJ*, 122, 2177
 Bauer, F. E. et al. 2002, *ApJ*, 123, 1163
 Boldt, E. 1987, *Phys. Rep.*, 146, 215
 Bolzonella, M., Miralles, J.-M., & Pelló, R. 2000, *A&A*, 363, 467
 Brandt, W. N., et al. 2001a, *AJ*, 122, 1
 Brandt, W. N., Hornschemeier, A. E., Schneider, D. P., Alexander, D. M., Bauer, F. E., Garmire, G. P., & Vignali, C. 2001b, *ApJ*, 558, L5
 Brandt, W. N., et al. 2001c, *AJ*, 122, 2810 (B01)
 Bruzual, A. G. & Charlot, S. 1993, *ApJ*, 405, 538
 Campana, S., Moretti, A., Lazzati, D., & Tagliaferri, G. 2001, *ApJ*, 560, L19
 Cohen, J. G., Hogg, D. W., Blandford, R., Cowie, L. L., Hu, E., Songaila, A., Shopbell, P., & Richberg, K. 2000, *ApJ*, 538, 29
 Coleman, G. D., Wu, C.-C., & Weedman, D. W. 1980, *ApJS*, 43, 393
 Cowie, L. L., Songaila, A., Hu, E. M., & Cohen, J. G. 1996, *AJ*, 112, 839
 Cowie, L. L., et al. 2001, *ApJ*, 551, L9
 Cowie, L. L., Garmire, G. P., Bautz, M. W., Barger, A. J., Brandt, W. N., & Hornschemeier, A. E. 2002, *ApJ*, 566, L5
 Crawford, C. S., Fabian, A. C., Gandhi, P., Wilman, R. J., & Johnstone, R. M. 2001, *MNRAS*, 324, 427
 Dawson, S., Stern, D., Bunker, A., Spinrad, H., & Dey, A. 2001, *AJ*, 122, 598
 Fabian, A. C. & Barcons, X. 1992, *ARA&A*, 30, 429
 Fabian, A. C. & Iwasawa, K. 1999, *MNRAS*, 303, L34
 Fabian, A. C., Wilman, R. J., & Crawford, C. S. 2002, *MNRAS*, 329, L18
 Fernández-Soto, A., Lanzetta, K. M., & Yahil, A. 1999, *ApJ*, 523, 72
 Fiore, F., et al. 2001, in *Issues of Unifications of AGNs*, ed. R. Maiolino, A. Marconi, & N. Nagar, in press (astro-ph/0109116)
 Gardner J. P. 1992, PhD Thesis, University of Hawaii.
 Gehrels, N. 1986, *ApJ*, 303, 336
 Giacconi, R., et al. 2002, *ApJS*, 139, 369
 Gruber, D. E., Rothschild, R. E., Matteson, J. L., & Kinzer, R. L. 1984, *Max Planck Inst. Rep.*, 184, 129
 Gruber, D. E. 1992, in *The X-ray Background*, ed. X. Barcons & A. C. Fabian, (Cambridge: Cambridge Univ. Press), p44
 Haiman, Z. & Loeb, A. 1999, *ApJ*, 521, L9
 Hasinger, G., Burg, R., Giacconi, R., Hartner, G., Schmidt, M., Trumper, J., & Zamorani, G. 1998, *A&A*, 329, 482
 Hasinger, G., et al. 2001, *A&A*, 365, L45
 Hasinger, G. 2002, in *New Visions of the X-ray Universe in the XMM-Newton and Chandra Era*, ed. F. Jansen, in press (astro-ph/0202430)
 Hawarden, T. G., Leggett, S. K., Letawsky, M. B., Ballantyne, D. R., & Casali, M. M. 2001, *MNRAS*, 325, 563
 Hodapp, K.-W. et al. 1996, *NewA*, 1, 177
 Hornschemeier, A. E., et al. 2001, *ApJ*, 554, 742 (H01)
 Hornschemeier, A. E., Brandt, W. N., Alexander, D. M., Bauer, F. E., Garmire, G. P., Schneider, D. P., Bautz, M. W., & Chartas, G. 2002, *ApJ*, 568, 82
 Hughes, D.H. et al. 1998, *Nature*, 394, 241
 Jenness, T. & Lightfoot, J. F. 1998, *Starlink User Note* 216.3
 Kron, R.G. 1980, *ApJS*, 43, 305
 Lehmann, I., et al. 2001, *A&A*, 371, 833
 Lightfoot, J. F., Jenness, T., Holland, W. S., & Gear, W. K. 1998, *SCUBA System Note* 1.2
 Loveday, J., Peterson, B. A., Efstathiou, G., & Maddox, S. J. 1992, *ApJ*, 390, 338
 Miyazaki, S., Sekiguchi, M., Imai, K., Okada, N., Nakata, F., & Komiyama, Y. 1998, in *Proc. SPIE Vol. 3355, Optical Astronomical Instrumentation*, ed. S. D’Odorico, 363-374.
 Moran, E. C., Lehnert, M. D., & Helfand, D. J. 1999, *ApJ*, 526, 649
 Morrison, R. & McCammon, D. 1983, *ApJ*, 270, 119
 Mushotzky, R. F., Cowie, L. L., Barger, A. J., & Arnaud, K. A. 2000, *Nature*, 404, 459
 Nandra, K., Mushotzky, R. F., Arnaud, K. A., Steidel, C. C., Gardner, J. P., Teplitz, H. I., & Windhorst, R. A., *ApJ*, in press (astro-ph/0205215)
 Norman, C., et al. 2002, *ApJ*, 571, 218
 Oke, J.B., et al. 1995, *PASP*, 107, 375
 Richards, E. A. 2000, *ApJ*, 533, 611
 Rosati, P., et al. 2002, *ApJ*, 566, 667
 Schmidt, M., et al. 1998, *A&A*, 329, 495
 Stern, D., et al. 2002a, *ApJ*, 568, 71

Stern, D., et al. 2002b, AJ, 123, 2223

Stocke, J. T., Morris, S. L., Gioia, I. M., Maccacaro, T., Schild, R.,

Wolter, A., Fleming, T. A., & Henry, J. P. 1991, ApJS, 76, 813

Tozzi, P., et al. 2001, ApJ, 562, 42

Vecchi, A., Molendi, S., Guainazzi, M., Fiore, F., & Parmar, A.N.
1999, A&A, 349, L73

Zezas, A., Georgantopoulos, I., & Ward, M. J. 1998, MNRAS, 301,
915




FIG. 1.— Schematic diagram of the 370 X-ray point sources (squares) in the B01 catalog of the CDF-N. Spectroscopically identified X-ray sources are solid. Radii of $6.5'$ and $10'$ from the approximate X-ray image center are indicated by the large circles. The dashed (solid) contour shows the area coverage of the HK' (submillimeter) observations. The optical data cover a larger field ($24' \times 24'$) than that shown.

FIG. 2.— z' -band thumbnail images of the 370 X-ray point sources in the B01 catalog. The 1σ AB magnitude limit is 27.0. The thumbnails are $18''$ on a side and are labeled with the B01 catalog number (column 1 of Table 1 in this paper) in the upper left corner. The expected position of each source is shown with a $2''$ radius circle if it is within $6.5'$ of the approximate X-ray image center and with a $3.6''$ radius circle otherwise. If an optically identified source is within $2''$ of the nominal X-ray position, then the spectroscopic redshift is shown in the lower left corner. If a bright optically identified source is within $5''$ of the nominal X-ray position, then the redshift of that source is printed with fainter text. Sources detected in the B01 2–8 keV catalog have an ‘H’ printed in the lower right corner. Sources also detected at 20 cm or $850\ \mu\text{m}$ have, respectively, an ‘R’ or an ‘S’ printed in the upper right corner. North is up and East is to the left.

FIG. 6.— Spectra for 175 of the 182 X-ray sources with spectroscopic redshifts versus rest-frame wavelength. The hatched regions show the positions of strong atmospheric bands and night sky emission lines. In each case we give the redshift and the B01 number. Sources with WIYN spectra are marked with a ‘W’ and are not flux calibrated.

TABLE 1. Optical, Submillimeter, and Radio Properties of the CDF-N X-ray Sample

B01	RA (2000)	Dec (2000)	850 μ m (mJy)	20 cm (μ Jy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
1	188.813873	62.23525	...	...	...	...	26.0	28.4	26.8	28.3	27.5	...	...	...	
2	188.819000	62.26025	...	...	...	...	-25.8	-25.3	-26.5	-25.7	-27.6	...	...	...	
3	188.838242	62.27450	...	...	21.8	...	21.1	20.8	21.7	22.2	22.9	0.8	0.0	0.559	
4	188.839798	62.25064	...	...	...	...	-26.9	26.6	27.7	27.7	27.2	...	...	...	
5	188.843094	62.23125	...	...	...	...	24.0	23.7	24.1	24.7	25.7	...	...	...	
6	188.854111	62.25658	...	...	...	...	24.6	24.8	25.6	26.6	26.4	...	...	...	
7	188.869492	62.24128	...	...	...	...	24.7	26.0	27.3	27.4	-30.8	...	...	...	
8	188.872330	62.21553	...	135 ± 11	23.1	...	22.0	22.0	22.8	23.4	23.8	0.0	0.8	obs	C
9	188.873505	62.30636	...	...	...	...	-26.7	26.9	26.6	29.3	28.4	...	...	...	
10	188.896500	62.24150	...	...	...	...	24.1	23.9	24.2	24.3	24.5	...	...	obs	
11	188.900421	62.31756	...	...	...	...	23.3	23.6	24.8	25.9	25.8	0.8	-0.8	...	
12	188.900711	62.27953	...	...	[18.0]	...	[18.0]	[18.0]	[18.0]	[18.0]	17.5	0.0	0.9	star	
13	188.905121	62.28986	...	...	22.3	...	21.9	21.5	22.2	22.1	22.4	-0.8	0.4	2.050	B
14	188.910461	62.27872	...	3550 ± 179	20.0	...	19.6	18.8	19.8	20.0	20.3	0.0	0.0	0.712	BC
15	188.912888	62.26683	...	...	...	...	23.3	22.9	23.1	23.1	23.4	0.4	-0.4	2.574	BC
16	188.913681	62.21994	...	...	...	...	27.3	-30.4	28.6	28.3	-28.1	...	...	...	
17	188.925949	62.32978	...	...	...	...	25.6	25.8	27.6	27.8	28.7	...	...	...	
18	188.928955	62.19553	...	...	...	...	26.0	25.3	25.8	27.3	26.9	...	...	...	
19	188.936127	62.29322	...	...	...	...	24.7	24.2	24.9	25.1	25.2	...	...	...	
20	188.939499	62.30006	...	...	...	-21.0	26.3	26.4	-29.6	-27.3	-30.9	...	...	...	
21	188.941254	62.23058	...	...	...	20.9	24.7	24.9	25.5	26.1	26.5	...	...	obs	
22	188.941803	62.26669	...	...	...	-21.0	25.2	25.1	28.0	28.5	28.5	...	...	...	
23	188.942459	62.29161	...	...	...	-20.6	-27.4	-25.9	-28.7	-25.8	-29.2	...	...	...	
24	188.948425	62.25375	...	...	...	21.6	24.6	24.7	25.5	25.7	26.0	...	...	...	
25	188.951248	62.28442	...	...	...	19.8	22.6	22.5	23.2	23.7	23.9	0.8	-0.4	obs	
26	188.951950	62.32542	...	...	...	...	25.2	24.9	25.3	26.2	27.4	...	...	...	
27	188.952164	62.25561	...	...	22.9	20.1	21.9	21.7	22.9	23.4	24.2	1.6	-0.4	0.885	
28	188.955826	62.26022	...	74 ± 9	...	19.3	22.8	22.6	23.1	23.4	24.0	0.0	0.0	obs	
29	188.960037	62.30252	...	...	...	-21.6	24.3	24.4	25.4	25.6	26.3	...	...	...	
30	188.960831	62.24875	...	...	20.1	18.5	22.5	22.2	22.7	23.7	24.8	0.0	0.0	obs	
31	188.965576	62.29925	...	...	...	-20.5	25.4	27.3	26.7	-27.9	27.7	...	...	...	
32	188.971451	62.17708	...	...	21.4	18.7	21.2	21.0	21.3	21.8	22.0	0.0	0.0	1.379	BC
33	188.974045	62.36533	...	83 ± 11	30.0	30.0	30.0	30.0	30.0	30.0	30.0	...	...	...	
34	188.975662	62.33664	...	...	30.0	30.0	30.0	30.0	30.0	30.0	30.0	...	...	...	
35	188.979218	62.26986	...	...	...	20.9	22.8	22.9	24.2	26.1	26.3	0.4	-0.8	1.021	
36	188.979752	62.30003	...	...	20.1	16.7	[18.0]	[18.0]	20.0	21.0	22.5	0.4	0.6	star	C
37	188.980545	62.25133	...	...	19.0	17.7	19.5	19.1	19.6	20.1	20.9	-0.4	0.4	0.207	
38	188.983414	62.26553	...	89 ± 9	21.6	18.3	20.6	20.5	21.4	22.5	23.9	0.4	0.0	0.436	C
39	188.983841	62.20530	...	...	...	18.8	22.0	22.1	23.7	24.7	25.9	0.0	0.4	obs	
40	188.984589	62.27145	...	...	18.4	16.2	18.1	[18.0]	18.8	19.7	21.2	0.0	0.6	0.296	
41	188.989426	62.25461	...	...	...	22.1	23.7	23.3	23.8	24.0	24.3	0.0	0.0	2.423	
42	188.990326	62.17350	...	...	22.6	20.7	22.4	21.9	22.2	22.4	23.2	-0.4	-0.4	3.067	BC

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μ m (mJy)	20 cm (μ Jy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
43	188.991257	62.21950	...	...	...	20.2	25.1	24.7	25.3	25.3	25.7	...	...	...	
44	188.991577	62.35342	...	...	...	...	22.0	22.0	23.1	23.8	24.3	0.0	-1.2	...	C
45	188.993896	62.18383	...	...	21.3	18.1	20.4	20.3	21.2	22.2	23.0	-0.4	0.8	0.637	
46	188.995499	62.26194	...	...	20.3	18.3	20.2	19.9	20.4	20.7	21.3	-0.8	-1.2	0.108	
47	188.998795	62.26386	...	212 $\pm$ 13	19.2	17.0	19.1	19.0	19.6	20.2	21.2	0.0	0.0	0.375	
48	189.000381	62.30453	...	...	...	20.5	22.5	22.6	23.7	24.3	24.6	0.4	-0.4	1.142	
49	189.001876	62.32375	...	...	...	...	23.6	23.1	23.4	23.6	24.3	0.0	-0.4	3.190	BC
50	189.006454	62.23500	...	...	...	-22.7	23.2	23.0	23.5	23.8	24.1	-0.4	0.8	obs	
51	189.012832	62.39180	...	...	19.8	...	19.2	19.2	20.1	20.9	22.1	-0.4	-0.4	0.333	
52	189.013870	62.18631	...	124 $\pm$ 10	20.8	17.6	19.2	19.8	20.6	21.7	22.4	-1.0	0.2	0.639	C
53	189.020385	62.33736	...	...	[18.0]	...	[18.0]	[18.0]	[18.0]	19.5	20.8	0.0	0.0	star	C
54	189.022506	62.33633	...	...	...	...	22.6	22.8	23.9	24.6	25.2	0.0	0.0	obs	
55	189.023910	62.16778	...	...	23.0	18.9	21.7	21.6	22.7	23.4	24.2	0.0	-1.2	0.936	
56	189.024047	62.19661	...	...	...	19.2	22.5	22.7	23.1	23.9	23.8	0.4	0.4	obs	C
57	189.024368	62.14386	...	...	...	21.2	26.0	25.2	26.6	26.6	-28.2	...	...	...	
58	189.026794	62.20895	...	...	...	21.1	24.8	24.9	25.9	26.4	27.4	...	...	...	
59	189.027496	62.16422	...	196 $\pm$ 12	22.1	18.3	21.2	21.2	22.2	23.5	24.7	0.0	0.0	obs	
60	189.027984	62.26408	4.4 $\pm$ 1.7	...	...	20.8	22.9	22.7	23.2	23.2	23.4	0.0	0.4	2.415	C
61	189.028213	62.20581	...	...	...	19.3	22.6	22.5	23.7	24.7	25.7	0.0	0.0	0.747	
62	189.031997	62.14739	...	...	20.4	19.7	22.3	22.3	22.8	24.1	25.7	0.0	0.0	obs	
63	189.032135	62.25095	1.0 $\pm$ 1.5	...	...	19.6	23.4	23.3	24.1	24.4	24.8	0.0	0.0	1.675	C
64	189.033950	62.17667	...	217 $\pm$ 13	21.6	18.5	20.7	20.4	21.6	22.8	23.8	-0.4	-0.4	0.681	
65	189.034378	62.26475	4.2 $\pm$ 1.4	59 $\pm$ 8	21.5	19.5	21.3	21.1	21.7	22.4	23.3	-1.6	0.8	0.459	
66	189.035873	62.24331	3.8 $\pm$ 2.1	68 $\pm$ 8	...	-23.1	24.2	23.5	23.9	24.5	24.9	...	...	...	
67	189.037262	62.30605	...	...	...	21.6	25.4	26.4	27.8	28.3	27.6	...	...	...	
68	189.040771	62.19622	...	...	18.8	17.1	18.7	18.6	19.2	19.8	20.8	0.8	-0.2	0.136	
69	189.046249	62.37456	...	154 $\pm$ 12	21.7	...	20.7	20.6	21.7	22.1	22.4	0.0	0.4	0.982	C
70	189.047027	62.15108	...	...	...	19.4	22.4	22.4	23.5	24.2	24.7	-0.4	-0.8	1.140	
71	189.047622	62.36461	...	...	22.9	...	22.8	22.4	22.8	23.2	24.1	0.0	0.0	0.294	
72	189.049346	62.17075	-4.9 $\pm$ 3.0	...	...	20.4	24.7	24.7	25.8	26.0	26.5	...	...	...	
73	189.049865	62.14997	...	...	...	22.1	23.7	23.6	24.9	25.3	25.7	-0.4	1.6	obs	
74	189.050079	62.19417	...	...	19.8	16.9	18.7	18.9	19.5	20.2	21.3	0.2	0.2	0.275	
75	189.050293	62.32800	...	...	21.1	...	20.7	20.6	21.1	21.1	21.1	0.0	0.8	2.217	BC
76	189.053878	62.32494	...	...	22.8	...	22.9	22.4	22.8	23.5	24.6	0.0	-0.8	3.938	BC
77	189.054489	62.20669	...	...	...	-21.7	28.0	26.1	26.6	27.2	26.8	...	...	...	
78	189.058837	62.17150	-2.4 $\pm$ 2.3	...	...	22.2	26.9	-28.3	-27.8	-27.1	-29.2	...	...	...	
79	189.060211	62.17933	0.8 $\pm$ 2.3	...	...	21.7	24.8	24.4	25.0	25.6	25.6	...	...	obs	C
80	189.060272	62.22189	...	...	20.7	17.6	20.0	19.9	20.7	21.7	22.7	0.4	0.4	0.454	
81	189.060623	62.12161	...	...	22.3	18.6	21.4	21.4	22.4	23.0	23.9	-1.2	0.8	obs	
82	189.062286	62.17411	-1.6 $\pm$ 2.2	...	21.2	17.3	18.9	19.0	21.0	22.0	23.3	1.4	-1.0	star	C
83	189.066207	62.25431	0.7 $\pm$ 1.1	...	...	21.4	26.6	26.3	28.2	26.9	-28.4	...	...	...	
84	189.066757	62.18553	-0.3 $\pm$ 2.3	...	...	20.6	24.4	24.4	25.4	26.3	26.2	...	...	...	

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μm (mJy)	20 cm (μJy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
85	189.067123	62.25375	5.7 ± 1.1	53 ± 8	...	21.9	24.6	24.3	25.3	26.2	27.3	...	...	...	
86	189.071044	62.16980	1.0 ± 2.0	62 ± 8	22.5	18.8	21.5	21.6	22.9	23.8	24.5	0.0	0.4	0.845	
87	189.073501	62.22892	...	...	21.6	[18.0]	20.6	20.5	21.5	22.8	24.3	0.4	0.4	0.534	
88	189.075088	62.27650	0.9 ± 1.2	47 ± 8	21.1	18.0	20.3	20.1	21.0	21.6	22.3	0.0	-0.4	0.679	C
89	189.077468	62.18750	1.5 ± 2.1	...	21.4	18.3	20.9	20.8	21.3	21.7	21.9	0.0	0.0	1.022	B
90	189.079910	62.24489	1.4 ± 1.5	...	...	21.2	25.6	24.5	25.3	26.4	27.3	...	...	...	
91	189.079879	62.40544	...	...	...	...	24.1	24.0	24.9	25.7	25.9	...	...	...	
92	189.081451	62.21466	...	108 ± 9	20.7	17.6	20.1	19.9	20.7	21.8	22.9	-0.4	0.0	0.473	
93	189.083038	62.31972	...	...	...	...	23.0	22.7	23.3	24.0	25.3	0.0	0.0	...	
94	189.085464	62.21058	...	...	...	20.2	24.0	24.6	25.7	26.3	26.7	...	...	obs	
95	189.086959	62.23756	2.4 ± 2.4	...	...	20.5	23.9	23.8	24.4	24.9	25.4	0.0	0.0	obs	
96	189.087585	62.23669	2.9 ± 2.4	...	...	20.5	23.8	23.6	24.4	25.2	25.6	0.0	0.4	obs	
97	189.087951	62.21767	6.5 ± 3.2	...	...	21.3	25.1	24.7	25.1	26.2	26.7	...	...	obs	
98	189.088409	62.18578	1.0 ± 1.8	52 ± 8	...	18.7	21.7	21.9	23.4	24.6	25.5	0.0	0.0	1.014	
99	189.091369	62.26756	0.6 ± 1.1	...	...	21.3	26.2	-28.4	27.0	29.0	28.2	...	...	...	
100	189.094406	62.17458	-0.1 ± 1.3	...	...	22.5	24.9	25.0	25.4	26.2	26.5	...	...	...	
101	189.095626	62.25744	-2.1 ± 1.1	...	20.5	18.6	20.2	20.0	20.3	20.3	20.5	0.0	0.0	2.590	BC
102	189.096069	62.22958	2.9 ± 2.9	...	21.3	18.3	20.7	20.5	21.2	22.0	22.8	-0.4	0.0	0.483	
103	189.098571	62.31044	...	...	...	19.3	21.5	21.7	23.0	23.9	24.9	0.0	-0.4	1.012	
104	189.098709	62.16917	-0.3 ± 1.3	...	...	21.1	-28.1	27.0	27.6	-27.2	-27.7	...	...	...	
105	189.102325	62.18650	1.5 ± 1.6	...	22.9	18.7	21.4	21.3	22.7	23.7	25.0	0.0	0.0	0.747	C
106	189.104416	62.35456	...	...	[18.0]	...	[18.0]	[18.0]	[18.0]	[18.0]	[18.0]	-0.3	-0.4	star	
107	189.105667	62.29381	0.6 ± 3.0	...	...	20.4	24.2	24.4	25.3	26.2	26.6	...	...	...	
108	189.105865	62.23466	0.4 ± 2.4	...	[18.0]	14.4	[18.0]	[18.0]	[18.0]	[18.0]	20.3	0.0	0.0	star	C
109	189.113678	62.21897	-2.0 ± 2.2	...	...	-23.1	-27.8	26.1	27.2	-29.0	27.7	...	...	...	
110	189.113831	62.21614	-3.6 ± 2.2	...	22.8	19.6	22.0	21.9	22.7	23.2	23.7	0.0	0.0	1.218	
111	189.114502	62.17389	1.6 ± 1.1	...	...	20.0	22.6	22.6	23.7	24.8	25.8	0.0	0.0	0.762	
112	189.114746	62.20506	1.0 ± 2.7	...	...	-22.5	-29.6	27.7	-27.2	-29.6	-28.4	...	...	...	
113	189.115677	62.19956	1.2 ± 2.3	...	...	22.1	23.6	23.3	24.0	24.8	25.9	-0.8	0.4	obs	
114	189.119171	62.36086	...	...	...	...	23.9	24.0	25.1	25.6	25.9	0.4	0.4	...	
115	189.119873	62.13511	...	...	...	20.3	25.0	24.9	25.4	26.3	26.8	...	...	...	
116	189.120377	62.10447	...	58 ± 9	22.2	18.8	21.4	21.6	22.4	23.0	23.5	0.0	0.4	1.264	
117	189.121338	62.17942	5.0 ± 1.2	81 ± 8	...	18.8	22.2	22.4	23.8	24.8	26.0	0.0	0.0	1.013	
118	189.122253	62.27050	1.0 ± 1.6	...	21.9	18.4	21.0	20.9	22.3	23.2	24.0	-0.4	0.4	0.847	
119	189.122711	62.25364	1.0 ± 1.7	...	...	21.2	23.3	23.0	23.7	24.3	25.4	0.0	0.0	obs	C
120	189.123001	62.37050	...	...	22.1	...	21.3	21.2	22.3	22.6	23.3	0.0	1.6	0.853	B
121	189.125244	62.15659	0.2 ± 1.6	46 ± 8	...	20.0	22.4	22.4	24.0	25.0	25.6	-0.4	0.4	0.953	
122	189.125214	62.09522	...	82 ± 9	21.0	[18.0]	20.1	20.0	20.8	21.7	22.6	-1.2	0.0	0.485	C
123	189.125885	62.43911	...	...	...	...	22.8	22.8	23.8	24.2	24.5	-0.4	-0.8	...	
124	189.130295	62.16619	4.0 ± 1.2	152 ± 10	...	21.0	25.0	25.0	25.7	26.0	27.4	...	...	...	
125	189.132004	62.15216	-0.8 ± 1.6	...	19.7	16.9	19.8	19.8	20.8	21.4	22.4	0.8	-0.8	0.845	
126	189.133041	62.29561	...	...	22.6	19.2	21.6	21.5	22.6	23.7	24.8	-0.4	1.2	0.680	

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μ m (mJy)	20 cm (μ Jy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
127	189.135117	62.13325	...	90 $\pm$ 9	...	20.9	22.9	22.9	23.7	24.0	24.1	0.8	0.4	1.995	
128	189.136551	62.34114	...	...	[18.0]	...	[18.0]	[18.0]	[18.0]	[18.0]	19.1	0.3	-1.1	star	
129	189.138382	62.14286	0.8 $\pm$ 2.3	...	22.0	18.4	21.0	21.0	21.8	22.2	22.6	0.0	0.4	0.934	C
130	189.139526	62.23836	2.1 $\pm$ 2.3	...	...	21.3	26.0	25.1	25.0	25.3	26.4	...	...	3.403	BC
131	189.140335	62.16825	1.1 $\pm$ 1.3	46 $\pm$ 8	22.6	18.8	21.6	21.6	22.8	23.5	24.3	-0.4	0.4	1.016	
132	189.140671	62.22050	-1.9 $\pm$ 1.7	...	...	21.3	25.3	25.7	25.6	26.3	26.8	...	...	...	
133	189.141083	62.22436	-0.9 $\pm$ 1.8	...	...	22.7	24.3	23.8	24.5	25.3	25.9	...	...	obs	
134	189.143600	62.20359	0.4 $\pm$ 2.2	233 $\pm$ 13	19.5	16.4	19.3	19.2	19.9	20.9	22.1	0.0	0.4	0.456	
135	189.143600	62.16142	4.0 $\pm$ 1.4	...	...	22.1	27.6	25.4	27.0	29.0	28.4	...	...	...	
136	189.143799	62.21156	2.4 $\pm$ 1.9	230 $\pm$ 13	...	19.1	22.1	22.3	23.3	23.9	24.3	0.0	-0.4	1.219	
137	189.143997	62.32389	...	98 $\pm$ 9	22.7	18.7	21.2	21.3	22.7	23.7	24.6	0.8	0.0	0.871	
138	189.144257	62.07744	...	...	...	...	24.9	24.8	26.4	28.4	27.4	...	...	obs	
139	189.145416	62.27450	-0.7 $\pm$ 3.2	...	...	18.9	22.1	22.0	23.3	24.3	25.1	0.0	0.4	0.847	
140	189.146957	62.19770	-0.6 $\pm$ 2.0	...	...	20.6	23.8	24.3	25.4	26.0	26.9	0.0	0.4	...	
141	189.147079	62.18608	2.1 $\pm$ 1.8	...	21.4	18.0	20.5	20.4	21.3	22.4	23.8	0.0	0.0	0.410	
142	189.148331	62.24011	3.0 $\pm$ 2.4	87 $\pm$ 8	...	19.3	23.0	23.0	23.6	23.9	24.2	0.0	-0.4	2.005	C
143	189.148453	62.11900	...	...	...	20.5	22.8	23.1	24.4	26.3	26.8	0.0	0.0	...	
144	189.152679	62.22969	0.7 $\pm$ 1.7	...	21.3	18.9	21.1	20.9	21.5	22.0	22.6	0.0	0.0	0.957	
145	189.153122	62.19889	7.0 $\pm$ 2.1	...	22.4	19.1	21.5	21.4	22.2	22.9	23.5	0.0	0.0	0.555	C
146	189.153275	62.37431	...	...	...	...	28.0	26.1	27.1	30.4	-30.6	...	...	...	
147	189.153656	62.22228	1.7 $\pm$ 1.6	50 $\pm$ 7	...	23.9	27.3	-26.4	27.5	-27.3	30.9	...	...	...	
148	189.154221	62.19303	0.7 $\pm$ 2.4	...	18.2	17.0	18.8	18.6	19.0	19.3	19.8	-0.8	0.0	0.081	
149	189.160461	62.22764	2.6 $\pm$ 1.3	...	...	19.0	22.1	22.3	23.3	24.3	25.2	0.0	0.0	0.357	
150	189.162354	62.17817	1.8 $\pm$ 1.5	...	...	22.2	26.3	-27.1	-27.3	-27.8	-28.5	...	...	...	
151	189.162796	62.16220	-1.0 $\pm$ 1.5	...	...	19.8	22.8	23.1	23.9	24.7	25.1	0.0	0.4	obs	
152	189.164917	62.20847	0.6 $\pm$ 1.1	...	...	20.6	24.1	23.8	24.5	25.4	26.0	...	...	obs	
153	189.165207	62.16019	-0.6 $\pm$ 1.6	...	...	20.8	-25.9	-29.7	29.0	-26.9	-29.0	...	...	...	
154	189.165375	62.16945	0. $\pm$ 1.5	...	22.2	19.2	21.4	21.3	22.1	23.1	24.0	-0.4	0.0	0.510	C
155	189.166672	62.21394	1.0 $\pm$ 1.1	...	22.2	18.5	21.1	21.0	22.1	22.8	23.7	-0.4	0.0	0.848	
156	189.167252	62.28217	...	...	...	19.0	21.5	21.5	23.1	24.2	25.4	0.4	0.0	0.943	C
157	189.167252	62.32800	...	...	[18.0]	12.6	[18.0]	[18.0]	[18.0]	[18.0]	18.8	0.0	0.7	star	
158	189.169937	62.17817	1.3 $\pm$ 1.7	...	...	22.2	25.0	24.6	25.6	25.4	26.1	...	...	obs	
159	189.173248	62.16336	-0.7 $\pm$ 1.8	75 $\pm$ 8	21.3	17.5	20.2	20.2	21.1	22.4	23.7	-0.4	0.0	0.518	
160	189.174164	62.19222	0.9 $\pm$ 2.2	...	19.9	18.3	20.0	19.7	20.1	20.3	20.9	-0.8	0.0	0.089	
161	189.175446	62.22544	1.5 $\pm$ 1.1	467 $\pm$ 24	...	21.8	25.3	24.9	25.7	26.1	28.2	...	...	...	
162	189.175888	62.28661	...	...	...	20.9	23.7	23.8	24.2	24.8	25.8	0.4	-0.4	obs	
163	189.175903	62.26272	-1.8 $\pm$ 3.0	150 $\pm$ 10	22.4	18.1	21.2	21.2	22.4	23.3	24.3	0.0	0.0	0.857	
164	189.176254	62.10336	...	...	...	-22.0	28.5	26.8	26.3	-29.3	29.3	...	...	...	
165	189.184799	62.19253	-0.3 $\pm$ 1.9	1290 $\pm$ 61	22.4	17.7	20.9	21.2	22.7	24.3	25.8	0.0	0.4	1.011	
166	189.187790	62.05244	...	...	22.4	...	22.1	21.8	22.2	22.2	22.5	0.4	-0.4	2.455	BC
167	189.189636	62.31403	...	...	...	22.3	24.7	24.6	25.3	25.4	25.7	...	...	...	C
168	189.191162	62.13169	...	48 $\pm$ 8	22.8	19.3	22.1	22.1	22.7	23.2	23.5	-1.6	0.4	1.433	

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μ m (mJy)	20 cm (μ Jy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
169	189.191711	62.36261	...	...	...	...	23.8	23.7	24.3	24.7	25.3	0.4	-0.4	obs	
170	189.191879	62.24694	10. $\pm$ 2.2	124 $\pm$ 9	...	-22.3	24.7	25.2	25.2	25.7	26.8	...	...	...	
171	189.193130	62.23466	0. $\pm$ 1.2	179 $\pm$ 11	22.3	18.2	20.9	21.0	22.2	22.8	23.8	-0.4	0.0	0.961	C
172	189.193375	62.25811	-0.3 $\pm$ 2.3	...	60.7	18.9	21.8	21.8	23.1	24.2	25.4	-0.4	-0.4	0.851	
173	189.194000	62.14925	0.9 $\pm$ 3.0	...	...	19.7	22.9	23.0	24.4	25.4	26.5	0.0	0.0	obs	
174	189.199707	62.16147	-4.9 $\pm$ 3.1	...	...	21.2	23.7	23.1	24.5	25.9	26.0	-0.8	-1.2	5.186	B
175	189.199799	62.17220	9.1 $\pm$ 3.1	...	...	21.0	26.3	25.7	-30.1	26.9	31.2	...	...	...	
176	189.200211	62.21919	0.4 $\pm$.82	...	21.1	17.9	20.3	20.2	21.1	22.3	23.7	0.4	0.4	0.475	
177	189.201248	62.24897	0. $\pm$ 2.1	...	...	-20.8	26.6	26.2	30.2	-27.2	29.3	...	...	...	
178	189.201584	62.24061	1.4 $\pm$ 2.0	...	19.3	17.1	19.0	18.6	19.2	19.5	20.1	-0.4	0.4	0.139	
179	189.203629	62.36550	...	...	...	...	25.2	-25.6	26.3	26.3	26.8	...	...	...	
180	189.203918	62.05700	...	...	22.7	...	21.2	21.3	22.7	23.7	24.5	-0.8	0.0	0.114	
181	189.204880	62.07756	...	850 $\pm$ 46	[18.0]	...	[18.0]	[18.0]	[18.0]	[18.0]	19.1	-1.6	0.0	0.113	
182	189.205627	62.31375	...	...	21.4	19.0	21.1	20.9	21.6	22.5	23.0	-0.8	-0.8	0.593	
183	189.206039	62.22978	-0.4 $\pm$.95	...	[18.0]	15.9	18.1	[18.0]	[18.0]	19.2	20.2	-0.1	-0.3	0.089	
184	189.206955	62.12720	...	307 $\pm$ 17	...	20.1	22.6	22.8	23.3	23.8	24.0	-0.8	0.8	2.315	
185	189.207336	62.22028	1.4 $\pm$.81	49 $\pm$ 7	22.4	18.8	21.4	21.3	22.0	22.9	23.9	0.0	0.4	0.475	
186	189.209763	62.36286	...	...	21.6	...	20.8	20.7	21.5	22.2	23.0	0.0	-1.2	0.476	C
187	189.210037	62.33484	...	...	...	...	22.6	22.7	24.2	25.2	26.5	-0.4	-0.8	1.011	
188	189.212952	62.17519	7.7 $\pm$ 4.7	95 $\pm$ 9	20.8	17.5	20.1	20.1	20.9	21.8	23.0	-0.4	0.4	0.410	
189	189.213547	62.18100	1.5 $\pm$ 3.4	...	...	19.8	22.9	23.5	24.8	26.3	27.0	0.0	0.0	obs	
190	189.215683	62.20595	4.5 $\pm$.80	49 $\pm$ 7	21.8	19.0	21.4	21.2	21.9	22.5	23.6	0.0	-0.8	0.401	
191	189.215958	62.25136	-2.5 $\pm$ 2.3	...	...	21.2	24.9	24.6	25.4	25.5	26.0	...	...	obs	
192	189.217377	62.46970	...	...	18.3	...	20.1	19.9	20.2	21.4	0.00	0.0	0.0	0.334	
193	189.220459	62.24561	1.7 $\pm$ 2.4	168 $\pm$ 11	19.6	16.6	19.1	18.9	19.8	20.7	22.1	0.0	0.4	0.321	
194	189.222534	62.19433	0. $\pm$ 1.2	65 $\pm$ 8	22.5	19.4	22.1	22.0	22.7	23.0	23.4	0.0	0.4	1.275	
195	189.223160	62.33858	...	...	...	...	23.1	23.2	24.3	25.4	26.0	0.0	0.0	1.023	C
196	189.223450	62.18764	3.9 $\pm$ 3.0	...	...	20.8	22.5	22.4	23.2	23.7	24.0	0.0	0.0	0.890	
197	189.227371	62.18639	-0.9 $\pm$ 2.7	...	...	21.3	25.5	26.0	25.9	26.7	27.0	...	...	...	
198	189.231094	62.21983	-0.7 $\pm$.91	...	...	18.8	21.6	21.9	23.3	24.3	25.5	0.0	0.4	0.955	
199	189.232498	62.20020	2.1 $\pm$.94	...	...	20.2	23.8	24.1	24.6	24.9	25.1	0.0	0.4	obs	
200	189.232742	62.13553	...	106 $\pm$ 9	22.5	17.8	21.1	21.1	22.5	23.7	25.0	0.4	0.8	0.792	
201	189.235748	62.25367	1.3 $\pm$ 2.8	...	...	22.0	-26.9	26.2	27.8	31.2	29.6	...	...	...	
202	189.236008	62.21261	-1.3 $\pm$.96	...	21.1	17.6	20.3	20.2	21.1	22.4	23.8	0.0	0.0	0.517	
203	189.237198	62.21711	-0.5 $\pm$.97	49 $\pm$ 7	...	19.3	22.4	22.6	23.6	24.2	24.8	0.0	0.0	0.474	
204	189.237076	62.08294	...	...	22.0	...	20.5	20.6	22.0	23.0	24.3	0.0	1.2	0.840	
205	189.238922	62.17355	-0.7 $\pm$ 2.5	...	...	19.8	22.4	22.3	23.6	24.4	25.0	0.0	1.2	0.845	C
206	189.239502	62.20280	2.2 $\pm$.99	...	21.7	17.9	20.8	20.7	21.9	23.0	24.1	0.0	0.8	0.663	
207	189.240005	62.05397	...	...	...	...	24.2	23.8	24.1	24.8	25.1	...	...	...	
208	189.241287	62.35780	...	...	...	...	23.5	23.6	24.1	25.2	25.7	0.0	0.0	...	
209	189.243362	62.16625	-0.1 $\pm$ 3.2	...	[18.0]	16.0	18.1	[18.0]	[18.0]	19.1	20.1	-0.5	-0.2	0.137	
210	189.244293	62.06758	...	...	21.5	...	21.4	21.2	21.4	21.4	21.6	0.4	-1.2	0.287	C

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μm (mJy)	20 cm (μJy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
211	189.245117	62.17286	-1.4 ± 2.4	...	...	20.1	24.5	24.2	25.2	25.4	25.9	...	...	obs	
212	189.245209	62.24303	-0.3 ± 2.2	...	21.6	17.9	20.6	20.4	21.5	22.2	23.0	0.0	0.4	0.678	
213	189.246674	62.30897	...	506 ± 30	...	22.2	-29.7	-28.7	26.2	27.2	27.3	...	...	...	
214	189.246628	62.42314	...	...	20.7	...	20.7	20.4	20.5	20.7	21.2	-1.6	-0.8	3.085	BC
215	189.249161	62.32595	...	...	...	20.5	22.6	22.8	23.8	24.6	25.0	0.0	1.2	1.144	
216	189.251923	62.25248	0.2 ± 4.1	...	...	22.0	24.0	23.8	24.4	24.8	25.5	...	...	obs	
217	189.254913	62.10770	...	...	19.9	16.5	18.8	19.1	20.2	21.4	22.6	1.2	0.4	0.561	
218	189.256256	62.31314	...	...	18.4	20.1	21.6	21.2	21.6	22.1	23.0	0.0	0.0	0.230	
219	189.257202	62.12239	...	...	...	-21.6	-25.6	-27.0	-27.4	-27.1	-27.8	...	...	...	
220	189.258484	62.18956	-1.1 ± 1.7	...	19.2	17.6	19.5	19.2	19.7	20.1	20.9	0.0	0.4	0.956	
221	189.259964	62.32383	...	...	...	21.9	25.9	25.4	25.5	26.0	25.7	...	...	...	
222	189.260834	62.21222	-1.7 ± 1.9	...	...	20.7	24.1	23.9	24.2	24.9	25.2	...	...	...	
223	189.261307	62.26214	...	...	19.9	17.2	19.5	19.4	20.0	20.5	21.0	0.4	0.4	0.514	B
224	189.261749	62.26703	...	...	...	21.6	24.6	24.9	25.4	26.7	26.2	...	...	obs	C
225	189.264832	62.27567	...	...	21.3	18.5	21.2	21.1	22.3	23.1	24.5	-1.6	0.8	0.359	
226	189.266998	62.19867	2.2 ± 1.7	...	...	-24.7	27.9	25.6	-28.6	26.7	30.9	...	...	...	
227	189.266998	62.13194	...	50 ± 8	22.0	18.4	21.4	21.4	21.9	22.3	22.8	0.0	0.8	1.253	BC
228	189.269196	62.28111	...	...	20.7	18.1	20.3	20.2	20.7	21.4	22.3	0.4	0.0	0.377	
229	189.269409	62.24139	2.0 ± 1.6	...	21.6	18.1	20.8	20.7	21.6	22.6	23.6	0.0	0.8	0.561	
230	189.270248	62.26711	...	...	...	21.0	25.2	24.9	25.8	27.4	27.0	...	...	...	
231	189.271255	62.27633	...	...	...	21.5	27.2	-26.6	27.9	28.6	-29.5	...	...	...	
232	189.274841	62.02530	...	...	22.2	...	21.0	20.9	22.0	23.2	24.5	0.8	-0.4	...	C
233	189.275879	62.36006	...	...	21.6	...	21.0	20.9	21.6	22.0	22.5	0.4	0.8	obs	C
234	189.277802	62.37433	...	...	...	...	28.5	-28.1	26.8	27.6	27.6	...	...	...	
235	189.277832	62.43058	...	...	20.1	...	19.5	19.5	20.1	20.8	21.1	0.4	0.0	...	C
236	189.278366	62.12294	...	72 ± 9	...	18.8	22.1	22.4	23.5	24.4	25.2	0.8	0.0	0.518	
237	189.278549	62.28397	...	...	19.8	18.0	19.2	18.8	19.5	19.6	20.0	0.4	0.0	1.018	BC
238	189.279404	62.14183	-2.8 ± 4.4	...	...	19.9	23.3	23.6	24.6	25.7	26.0	-0.4	-0.4	obs	
239	189.279785	62.27845	...	...	...	19.7	22.7	22.9	24.0	24.6	24.9	2.0	0.0	1.017	
240	189.280121	62.23555	6.2 ± 1.3	45 ± 7	...	20.3	24.4	24.6	25.4	25.9	26.9	...	...	...	
241	189.281113	62.33220	...	...	...	...	23.5	23.9	25.3	26.7	-29.6	1.2	0.0	...	
242	189.281158	62.36341	...	70 ± 10	22.4	...	22.0	22.0	22.3	23.0	23.1	0.4	-0.4	1.451	BC
243	189.281876	62.09278	...	...	...	...	26.8	25.0	26.3	27.2	27.7	...	...	...	
244	189.282288	62.27117	...	...	22.7	18.5	21.4	21.5	22.9	23.7	24.6	0.0	0.4	0.940	
245	189.283035	62.26828	...	...	22.5	18.5	21.2	21.2	22.5	23.2	24.0	-0.4	0.4	0.936	
246	189.284622	62.18220	-1.0 ± 1.6	45 ± 8	20.5	17.9	20.2	20.1	20.5	21.0	21.9	0.0	0.0	0.422	
247	189.286072	62.25042	0.8 ± 2.3	...	22.3	19.0	21.5	21.3	22.3	23.5	25.0	0.0	0.0	0.568	
248	189.287262	62.08845	...	...	22.7	...	21.2	21.4	22.9	24.1	25.4	-1.6	0.8	0.956	
249	189.288757	62.37058	...	...	...	...	23.6	23.5	24.9	25.6	26.2	0.8	0.8	...	
250	189.289368	62.14364	5.4 ± 4.5	72 ± 8	22.1	17.6	20.5	20.6	22.0	23.2	24.4	0.0	0.4	0.907	
251	189.290955	62.13356	...	...	23.2	19.8	22.2	22.2	22.9	22.9	23.1	0.0	0.4	2.182	BC
252	189.291077	62.14267	6.0 ± 4.8	...	...	19.6	22.6	22.8	24.3	26.3	-29.5	0.0	0.0	obs	

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μm (mJy)	20 cm (μJy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
253	189.291458	62.38308	...	708 $\pm$ 36	...	...	22.9	23.3	24.4	25.3	25.9	0.0	0.0	...	
254	189.292252	62.09647	...	...	...	-21.4	23.6	23.4	24.1	25.2	26.0	0.0	0.0	...	
255	189.295975	62.39097	...	...	...	...	23.1	23.2	23.6	24.3	24.5	0.4	0.8	1.392	BC
256	189.296539	62.12033	...	...	...	20.3	24.2	24.4	25.3	26.5	28.9	...	...	...	
257	189.297379	62.22528	0.7 $\pm$ 1.3	132 $\pm$ 10	...	19.8	23.1	22.9	23.7	24.1	24.8	-0.8	0.4	obs	
258	189.299164	62.26322	...	...	...	20.2	22.8	23.0	24.3	25.7	26.7	0.4	0.0	obs	
259	189.299866	62.22367	4.1 $\pm$ 1.3	53 $\pm$ 8	...	21.5	25.1	24.8	25.5	26.2	26.2	...	...	...	
260	189.300446	62.20314	8.0 $\pm$ 1.8	...	...	21.9	25.8	25.3	26.5	27.9	28.9	...	...	...	
261	189.300507	62.29833	...	...	...	19.3	22.9	23.1	23.9	24.6	25.4	0.4	0.0	obs	
262	189.302750	62.22847	0.3 $\pm$ 1.2	...	...	-22.6	26.1	-26.2	-27.5	27.5	29.4	...	...	...	
263	189.307083	62.26266	...	...	...	21.2	-26.1	25.5	26.9	27.6	27.8	...	...	...	
264	189.307541	62.24014	1.8 $\pm$ 1.6	...	22.2	18.6	21.3	21.2	22.1	23.1	24.3	0.0	0.4	0.475	
265	189.307877	62.24944	-1.0 $\pm$ 2.3	...	...	21.5	28.7	25.7	25.6	26.5	26.2	...	...	...	
266	189.307999	62.30744	...	595 $\pm$ 30	...	23.7	-25.4	25.1	28.5	-27.2	-27.3	...	...	...	
267	189.308624	62.15464	...	...	...	20.1	23.9	24.1	24.8	25.7	26.3	0.0	0.0	obs	
268	189.309586	62.20228	4.5 $\pm$ 1.9	...	...	20.6	23.3	23.4	23.7	24.0	24.5	0.0	0.8	...	
269	189.309708	62.20583	2.2 $\pm$ 2.2	...	...	20.5	23.0	23.0	24.2	25.0	25.5	0.4	0.8	1.084	C
270	189.310410	62.31083	...	...	...	21.5	24.9	24.5	25.3	25.8	26.8	...	...	...	
271	189.310791	62.08430	...	58 $\pm$ 9	...	...	22.6	22.4	23.0	23.9	24.9	0.0	0.0	...	
272	189.311462	62.27136	...	...	...	20.4	23.4	23.1	23.9	24.1	24.4	0.4	0.4	obs	
273	189.313492	62.04203	...	...	...	...	24.8	25.3	25.8	26.4	27.2	...	...	...	
274	189.316452	62.20364	1.9 $\pm$ 2.2	...	...	19.0	22.2	22.4	23.7	24.7	25.3	0.0	0.4	1.019	
275	189.316422	62.05661	...	109 $\pm$ 11	19.9	...	19.2	18.8	19.7	19.8	20.2	0.8	-0.4	2.070	BC
276	189.317459	62.11889	...	...	21.2	18.0	20.7	20.4	21.2	21.3	21.7	-0.4	0.4	0.679	BC
277	189.317627	62.19175	-0.2 $\pm$ 1.7	...	...	19.8	22.6	23.0	24.4	27.0	27.8	-0.4	0.4	obs	
278	189.318207	62.25356	...	187 $\pm$ 12	20.9	17.5	20.2	20.1	21.2	22.2	23.5	0.0	0.0	0.230	
279	189.318802	62.27867	...	66 $\pm$ 8	22.4	19.6	22.1	21.8	22.4	23.1	23.7	1.2	1.6	0.557	
280	189.319412	62.29261	...	346 $\pm$ 19	...	18.9	22.1	22.1	23.2	24.0	24.9	0.4	0.0	1.146	C
281	189.319946	62.10653	...	...	...	19.4	22.5	22.5	23.6	24.2	24.8	0.0	0.0	obs	
282	189.320038	62.09806	...	...	21.9	19.6	21.6	21.2	21.8	21.8	22.2	0.8	0.4	1.939	BC
283	189.324295	62.31547	...	...	22.2	21.4	21.8	21.6	22.1	22.1	22.3	0.4	0.4	2.218	BC
284	189.328506	62.11444	...	...	...	21.0	23.5	23.2	23.8	24.1	24.4	0.0	0.0	2.729	C
285	189.329300	62.17377	-0.4 $\pm$ 3.5	...	...	-21.8	25.3	25.0	25.7	-27.8	30.8	...	...	4.137	
286	189.332748	62.07583	...	...	...	...	23.1	23.2	24.2	25.2	25.7	0.0	0.0	...	
287	189.332916	62.16533	...	...	...	21.7	24.0	23.9	24.4	24.8	25.2	...	...	2.638	C
288	189.334702	62.23131	1.3 $\pm$ 1.7	...	...	19.4	22.6	22.6	23.9	24.9	25.9	0.0	0.8	obs	
289	189.335007	62.25653	...	...	...	18.5	21.5	21.8	23.3	24.9	26.3	0.0	0.0	0.935	
290	189.340179	62.21292	2.8 $\pm$ 2.7	...	17.5	17.4	19.5	19.9	20.2	20.2	21.5	0.0	0.0	0.104	
291	189.341125	62.17647	12. $\pm$ 4.1	...	22.6	18.5	21.3	21.3	22.6	23.3	24.1	0.0	0.4	0.978	
292	189.343246	62.09341	...	...	...	...	25.9	-27.2	27.9	-26.9	27.6	...	...	...	
293	189.344635	62.15978	...	...	...	22.2	24.8	24.2	24.6	24.9	25.2	...	...	obs	
294	189.346664	62.26069	...	...	...	21.4	23.7	23.5	24.1	24.4	24.8	0.0	0.4	2.240	C

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μ m (mJy)	20 cm (μ Jy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
295	189.348053	62.35353	1.1 ± 1.5	...	...	...	23.6	23.5	23.8	24.1	25.2	1.2	0.8	...	C
296	189.350082	62.21783	...	...	...	20.6	23.7	24.0	24.5	24.9	25.2	0.0	0.0	obs	
297	189.350708	62.21119	...	...	...	22.6	25.0	24.1	25.5	27.0	29.2	...	...	...	
298	189.351196	62.23320	...	...	...	20.5	24.1	23.9	24.9	25.3	25.7	...	...	obs	
299	189.354416	62.14906	...	90 ± 9	...	18.7	22.0	22.2	23.7	25.0	25.8	-0.4	0.0	obs	
300	189.356171	62.32811	-1.1 ± 1.4	122 ± 10	18.6	...	18.9	18.7	19.2	19.7	20.7	0.4	0.4	0.279	
301	189.356339	62.28544	...	...	...	21.5	25.7	26.4	26.1	-27.2	28.3	...	...	...	
302	189.357132	62.28017	...	...	20.0	16.0	[18.0]	[18.0]	19.7	20.7	22.1	0.0	0.0	star	C
303	189.357285	62.37883	...	...	...	...	22.9	22.7	23.2	23.1	23.5	-0.8	-0.8	2.214	C
304	189.357498	62.04192	...	...	21.1	...	20.5	20.4	21.1	21.5	22.0	0.4	0.4	0.605	C
305	189.360367	62.07592	...	...	...	...	22.9	22.9	23.5	23.9	24.3	0.0	0.0	obs	
306	189.360840	62.34119	-0.8 ± 1.3	102 ± 10	...	...	24.1	24.1	25.0	25.3	26.0	...	...	...	
307	189.361603	62.29655	...	...	...	21.3	24.3	24.9	25.0	25.3	25.7	...	...	obs	
308	189.374039	62.21019	...	...	22.5	20.5	22.6	22.2	22.6	22.7	23.4	-0.4	-0.4	0.159	
309	189.378540	62.14717	...	...	22.5	18.3	21.1	21.1	22.5	23.6	24.7	-0.4	-0.4	0.777	
310	189.379578	62.17522	...	...	...	21.7	26.2	25.0	26.6	27.5	28.6	...	...	obs	
311	189.381638	62.21811	...	...	...	21.4	24.7	24.5	25.3	25.7	26.7	...	...	...	
312	189.382233	62.28431	...	...	...	21.2	23.2	23.2	24.0	24.6	24.9	-1.2	0.0	...	
313	189.384125	62.21275	...	...	23.0	18.2	21.4	21.6	22.9	23.9	24.7	-0.4	0.8	1.022	
314	189.391998	62.27331	...	...	...	20.2	25.3	25.2	25.8	-30.5	-27.8	...	...	...	
315	189.392380	62.15873	...	142 ± 11	19.3	16.5	18.8	18.4	19.4	20.0	21.1	0.8	0.0	0.189	
316	189.393951	62.23220	4.3 ± 2.1	...	21.7	18.0	20.8	20.7	21.9	22.7	23.5	0.0	0.4	0.839	
317	189.395584	62.32114	-3.1 ± 2.8	...	...	20.8	23.7	23.5	24.1	24.6	25.4	0.0	0.0	...	
318	189.396378	62.21861	...	...	...	19.0	22.2	22.2	23.3	24.6	26.1	0.0	0.0	obs	
319	189.397705	62.23167	4.1 ± 2.0	...	...	19.8	23.5	23.5	24.4	25.2	25.6	-0.8	0.0	obs	
320	189.399841	62.30156	...	...	...	21.1	23.1	23.2	23.7	24.5	25.1	0.0	0.0	...	
321	189.402420	62.18089	...	...	22.1	20.6	22.3	22.1	22.3	22.5	22.8	-0.8	1.2	0.033	
322	189.403717	62.24136	1.7 ± 1.4	57 ± 8	...	20.4	24.7	26.6	26.2	26.6	27.0	...	...	...	
323	189.404541	62.30956	...	...	...	21.5	25.2	25.9	27.6	30.5	28.8	...	...	...	
324	189.407532	62.27508	...	...	[18.0]	13.9	[18.0]	[18.0]	[18.0]	[18.0]	[18.0]	0.3	0.3	star	
325	189.408386	62.28708	...	...	...	21.4	23.9	23.7	24.0	25.7	26.0	0.0	0.0	obs	C
326	189.408478	62.21911	6.1 ± 4.0	...	...	19.4	22.9	23.1	24.7	26.3	26.6	-0.8	0.4	1.238	
327	189.413483	62.34995	...	72 ± 10	20.9	...	20.2	20.1	20.8	20.9	21.5	0.4	-0.4	0.837	BC
328	189.415298	62.22703	1.9 ± 1.6	...	22.5	18.1	21.0	21.0	22.5	23.5	25.0	0.8	0.4	0.838	
329	189.415573	62.28378	...	...	21.1	17.9	20.7	20.5	21.4	22.4	23.7	0.0	0.0	0.458	
330	189.420746	62.20003	...	...	22.4	18.7	21.7	21.7	22.5	22.9	23.3	0.0	0.4	1.168	
331	189.421463	62.17984	...	...	...	20.1	25.0	24.6	27.1	26.7	27.1	...	...	...	
332	189.422470	62.21419	...	163 ± 11	...	19.0	22.5	22.5	23.6	24.2	24.8	-0.4	0.4	obs	
333	189.426376	62.25506	-0.9 ± 1.6	...	18.8	17.1	18.6	18.3	19.0	19.4	20.1	-0.2	0.2	0.069	
334	189.427673	62.30339	...	...	21.7	19.5	21.2	21.1	21.5	21.5	21.8	-0.8	-0.4	2.302	BC
335	189.428329	62.13131	...	...	22.3	18.8	21.5	21.5	22.3	22.9	23.5	-1.2	-1.2	1.084	
336	189.431747	62.09422	...	...	...	...	25.4	25.4	26.4	28.8	29.7	...	...	...	

TABLE 1. (continued)

B01	RA (2000)	Dec (2000)	850 μ m (mJy)	20 cm (μ Jy)	$R(iso)$	HK'	z'	I	R	V	B	Δx^a ($''$)	Δy^a ($''$)	z	type ^b
337	189.435577	62.11150	...	...	...	23.2	-28.8	-26.8	-27.3	-26.5	-28.4	...	...	...	
338	189.437119	62.12208	...	...	...	21.2	-26.1	-25.4	27.4	-28.0	30.0	...	...	...	
339	189.442703	62.17656	...	...	...	22.2	-25.7	25.9	25.9	27.3	27.0	...	...	obs	
340	189.444733	62.12047	...	...	...	21.0	25.2	25.4	26.8	25.9	27.5	...	...	...	
341	189.450378	62.35725	...	...	[18.0]	...	[18.0]	[18.0]	[18.0]	22.0	24.0	0.0	0.0	star	C
342	189.459137	62.23297	0.3 ± 1.3	90 ± 9	...	20.7	24.4	24.8	26.3	27.2	27.1	...	...	obs	
343	189.469589	62.27439	...	79 ± 9	19.2	16.6	18.7	18.5	19.3	19.9	20.5	0.2	0.6	0.307	B
344	189.469635	62.12228	...	86 ± 10	22.3	19.0	21.2	21.4	22.7	23.5	24.3	-0.8	-0.4	0.968	
345	189.476181	62.18317	...	285 ± 16	...	19.0	22.6	23.1	24.2	25.5	25.7	1.2	1.6	obs	
346	189.477219	62.16911	...	...	...	-20.9	25.4	24.2	25.5	26.8	26.9	...	...	obs	
347	189.477539	62.31175	...	...	...	...	26.6	25.0	25.7	26.1	26.6	...	...	...	
348	189.488876	62.27442	...	...	22.0	22.2	21.7	21.4	21.8	22.6	23.2	-0.4	-0.4	2.922	BC
349	189.498459	62.18391	...	85 ± 10	[18.0]	16.7	18.2	[18.0]	[18.0]	[18.0]	18.8	-0.5	0.6	0.909	B
350	189.503204	62.14436	...	...	...	-20.7	25.8	26.7	27.2	-27.6	28.1	...	...	...	
351	189.504028	62.22669	4.2 ± 2.0	190 ± 13	[18.0]	16.6	[18.0]	[18.0]	[18.0]	19.1	19.2	-0.4	-0.2	0.443	
352	189.513092	62.18453	...	...	...	19.7	23.3	23.3	23.9	24.6	25.0	0.0	0.0	obs	
353	189.518616	62.25686	...	...	...	-21.9	23.4	23.3	23.9	24.2	24.6	-1.2	1.2	obs	
354	189.521927	62.12750	...	...	22.3	...	21.3	21.3	22.3	22.9	23.7	-0.8	1.2	...	
355	189.525452	62.17970	...	...	20.3	17.1	19.8	19.7	20.5	21.4	22.2	-1.6	0.8	0.570	
356	189.528778	62.21197	-1.5 ± 1.9	1010 ± 53	22.4	...	21.0	21.3	22.8	24.3	25.2	-1.2	-0.8	1.024	
357	189.544052	62.29133	...	...	...	...	22.8	23.2	24.2	25.2	26.1	-0.4	0.8	...	
358	189.555740	62.14953	...	...	...	...	24.2	24.3	25.6	26.9	26.6	...	...	...	
359	189.556091	62.19280	-4.7 ± 3.9	...	...	...	27.2	-25.9	-28.4	-27.0	-30.9	...	...	...	
360	189.569946	62.18178	...	...	22.6	...	21.4	21.6	22.6	23.2	23.8	0.4	-1.2	obs	
361	189.572830	62.26914	...	...	...	...	22.8	22.9	23.3	24.0	24.0	-1.2	0.4	...	
362	189.593246	62.23797	...	...	22.8	...	21.6	21.7	22.8	23.5	24.0	-0.8	0.0	0.986	
363	189.596161	62.17747	...	...	...	...	24.4	24.5	26.2	26.7	29.7	...	...	...	
364	189.597916	62.16211	...	...	...	...	25.1	24.8	25.3	25.8	25.9	...	...	...	
365	189.599289	62.25472	...	...	...	...	25.8	30.9	25.7	27.4	27.3	...	...	...	
366	189.618332	62.15092	...	...	22.0	...	21.8	21.5	21.8	22.0	22.5	-1.6	1.2	...	C
367	189.630295	62.20628	...	...	22.7	...	22.3	21.9	22.5	22.5	23.0	-0.8	-0.8	...	C
368	189.642166	62.24383	...	...	...	...	26.3	-25.6	27.4	-26.6	-27.6	...	...	...	
369	189.670975	62.20042	...	...	22.5	...	21.5	21.6	22.8	23.4	24.0	-0.8	-0.4	...	
370	189.689728	62.16689	...	...	[18.0]	...	[18.0]	[18.0]	[18.0]	[18.0]	[18.0]	0.0	1.3	star	

^aPositive offsets are to the East and North.^bB indicates the spectrum has broad emission lines. C indicates the source is optically compact.